



High genotypic variation of cocktail tomato yield and quality under different potassium supply

Frederike Wenig¹, Marcel Naumann¹, Bernd Steingrobe², Elke Pawelzik¹ and Inga Smit^{1}*

¹Quality of Plant Products, Department of Crop Sciences, University of Göttingen, Carl-Sprengel-Weg 1, 37075 Göttingen;

²Plant Nutrition and Crop Physiology, Department of Crop Sciences, University of Göttingen, Carl-Sprengel-Weg 1, 37075 Göttingen
Email*: ismit@gwdg.de

There is evidence that potassium (K) fertilization enhances the yield of medium and large-sized tomato cultivars. Many authors observed positive effects of K fertilization on fruit color, texture, total soluble solids (TSS) and titratable acids (TA) [3, 6]. The market for small-sized cocktail tomatoes is currently growing due to their preferable taste [5] which is positively related to increasing levels of TSS and TA [1, 2, 4].

We hypothesize, that K improves yield and quality of cocktail tomatoes. In order to test our assumption, we carried out pot experiments in two consecutive years. In a pot (6 L) experiment, three cocktail tomato cultivars (Primavera, Resi and Yellow Submarine) were fertilized with five K levels (3.1, 6.3, 9.4, 12.5 and 18.7 g K/pot applied in total over 19 weeks) to find out ideal dosages for well and deficient fertilized plants. Interestingly, the K effect on yield and most quality parameters was highly cultivar specific: while Primavera and Yellow Submarine reached a yield increase of more than 100% from the lowest to the highest K dose, no response of Resi was observed. In contrast to Primavera and Yellow Submarine, Resi showed only weak K fertilization effects on fruit quality aspects. Therefore, in a second experiment we grew the cultivars Primavera and Resi with two K doses (deficient and well-fertilized). In a third treatment, we interrupted K fertilization in order to induce a response in the cv. Resi to a sudden shortage of K. In contrast to Primavera, Resi showed no yield response to K fertilization like in the first experiment. The low yielding cv. Resi seemed to be more K efficient compared to Primavera. We conclude that the effect of K on yield and quality of cocktail tomatoes depends on the cultivar.

Literatur

- [1] Beckles, D.M. (2012): Factors affecting the postharvest soluble solids and sugar content of tomato (*Solanum lycopersicum* L.) fruit. *Postharvest Biology and Technology*, 63, pp 129–140.

- [2] Fontes, P.C.R., Sampaio, R.A. and Finger, F.L. (2000): Fruit Size, Mineral Composition and Quality of Trickle-irrigated Tomatoes as affected by Potassium Rates. *Pesquisa Agropecuária Brasileira*, 35, pp 21 – 25.
- [3] Hartz, T.K., Johnstone, P.R., Francis, D.M. and Miyao, E.M. (2005): Processing Tomato Yield and Fruit Quality Improved with Potassium Fertigation. *HortScience*, 40 (6), pp 1862 – 1867.
- [4] Javaria, S., Khan, M.Q. and Bakhsh, I. (2012): Effect of Potassium on chemical and sensory attributes of tomato fruit. *The Journal of Animal & Plant Sciences*, 22 (4), pp 1081 – 1085.
- [5] Laber, H. and Lattauschke, G. (2014): *Gemüseanbau*. 2. Auflage, Verlag Eugen Ulmer, Stuttgart (Hohenheim)
- [6] Liu, K., Zhang, T.Q., Tan, C.S. and Astatkie, T. (2011): Responses of Fruit Yield and Quality of Processing Tomato to Drip-Irrigation and Fertilizers Phosphorus and Potassium, *Agronomy Journal*, 103 (5), pp 1339 – 1345.