

## Susceptibility of red flour beetle *Tribolium castaneum* (Herbst) populations from Serbia to contact insecticides

Andrić, G.\*#, Kljajić, P., Perić, I., Pražić Golić, M.

Pesticides and Environment Research Institute, Banatska 31b, 11080 Belgrade, Serbia.

Email: goran.andric@pestring.org.rs

\* Corresponding author

# Presenting author

DOI: 10.5073/jka.2010.425.103

### Abstract

Contact insecticides remain the principal option for controlling stored-product insects. Unsatisfactory results of insecticide applications are caused by several factors, one of the most important being resistance of stored-product insects. The objective of this study was to examine the susceptibility in several populations of red flour beetle *Tribolium castaneum* (Herbst) from Serbia to different contact insecticides. Toxicity of the insecticides dichlorvos, malathion, chlorpyrifos-methyl, pirimiphos-methyl, deltamethrin and bifenthrin to adults of a laboratory population of *T. castaneum* was investigated in the laboratory by topical application. At the LD<sub>50</sub>, deltamethrin was the most toxic and malathion the least toxic of the insecticides. Discriminating dose data for the laboratory population were used to test the susceptibility of 10 other populations originating from different storage facilities (silos, warehouses and flour mills) in Serbia. The discriminating dose of malathion caused mortality of up to 85% in seven populations, indicating malathion resistance in those populations. For two populations of *T. castaneum* from Nikinci and Jakovo LD values, *ld-p* lines and levels of susceptibility/resistance (RRs) were determined. The most toxic insecticide for adults from Nikinci and Jakovo was deltamethrin, while malathion was least toxic. The resistance ratios (RRs) for malathion at the LD<sub>50</sub> were 17.6 for beetles from Nikinci, and 26.0 for beetles from Jakovo.

Keywords: *Tribolium castaneum* adults; Different populations; Insecticide toxicity; Susceptibility resistance

### 1. Introduction

Red flour beetle *Tribolium castaneum* (Herbst) (Coleoptera: Tenebrionidae) is a cosmopolitan species and one of the most widespread insects in storage facilities worldwide, especially in mills, and it develops most rapidly under favourable conditions (Rees, 2004). In Serbia, *T. castaneum* is common in grain storages, mills and flour depots, as well as in foodstuff production plants (Almaši, 2008). Application of contact insecticides and fumigants continues to be the most significant tools for protection of stored products from insect pests. Contact insecticides ensure long-term protection and prevent infestation of treated products. Several formulations based on malathion, pirimiphos-methyl and synergized or nonsynergized deltamethrin have been registered in Serbia for treatment of grains, and dichlorvos for treatment of vacant storage areas (Kljajić, 2008).

The efficacy of contact insecticides in protecting stored products is affected by several factors, primarily by altered susceptibility or resistance of stored-product insects (Subramanyam and Hagstrum, 1996; Kljajić and Perić, 2005). According to data collected globally by the FAO organization (Champ and Dyte, 1976), susceptibility to malathion had changed in 87% of the examined populations of *T. castaneum*. In the USA, many populations collected from peanut, barley and wheat storages were resistant to malathion (Halliday et al., 1988; Subramanyam et al., 1989; Zettler and Cuperus, 1990). Populations of *T. castaneum* from mills were resistant to malathion, and also dichlorvos and chlorpyrifos-methyl (Zettler, 1991; Zettler and Arthur, 1997). For the purpose of monitoring *T. castaneum* resistance to insecticides in Serbian storage facilities, we determined the toxicity parameters (LD levels and *ld-p* lines) for a laboratory population of *T. castaneum* by topical application and determined levels of susceptibility/resistance (resistance ratios, RRs) in populations collected from different silos, warehouses and flour mills.

## 2. Materials and methods

### 2.1. Populations tested and contact insecticides used

A laboratory population of *T. castaneum* was used in the experiment, as well as populations collected in 2008 and 2009 from various storage facilities in Serbia: (1) Bačka Topola (silo and flour mill) and Čurug (silo) in the north of Serbian province of Vojvodina; Novo Miloševo (silo) and Kikinda (flour mill) in the northeast; and Adaševci (silo), Nikinci (warehouse) and Irig (silo) in the southwest part, and (2) Belgrade Port (silo) and Jakovo (silo) of Belgrade area.

All populations were reared using methods described by Harein and Soderstrom (1966) and Davis and Bry (1985). Two-to-four-week-old unsexed insects were used in all tests and  $F_3$  generation of beetles for the susceptibility tests of collected *T. castaneum* populations. Technical concentrates of the following insecticide active ingredients were used in bioassay: dichlorvos 98%, malathion 96%, chlorpyrifos-methyl 97%, pirimiphos-methyl (product Actellic 50 EC containing 50% a.i.), deltamethrin 98% and bifenthrin 94.7%.

### 2.2. Bioassay

Insecticide toxicity to *T. castaneum* adults and susceptibility/resistance of the collected populations were tested by topical application in the laboratory at  $24\pm 1^\circ\text{C}$  and  $60\pm 5\%$  r.h. (Halliday et al., 1988). To immobilize the insects, they were anesthetized with  $\text{CO}_2$  for about 30 seconds before  $0.5 \mu\text{L}$  of each insecticide, dissolved in acetone (6-8 concentrations), was applied to the thorax of each insect by Burkard microapplicator (needle No. 18 in 1.0 mL syringe). Control insects were treated with acetone alone. After treatment of 25 insects in four replications, the beetles were transferred to clean Petri dishes after 4-6 h, each was filled with approximately 1 g of wheat flour. Lethal effects were determined after 7 d of microapplication. Based on discriminating doses causing 100% mortality in the laboratory population and reactions of each collected population to them, the populations from Nikinci and Jakovo were chosen for a full-range testing of insecticide toxicity or susceptibility.

### 2.3. Data analysis

Mortality data for treated insects were corrected for mortality in the control using Abbott's (1925) formula. Data were processed by probit analysis according to a method described by Finney (1971) and using a computer software developed by Raymond (1985). Statistical significance of differences between toxicity indicators for the insecticides investigated was assessed based on the overlapping/non-overlapping of intervals of confidence.

## 3. Results

Data on the toxicity of contact insecticides to adults of the laboratory population of *T. castaneum* (Table 1) show that deltamethrin was the most toxic insecticide at the  $\text{LD}_{50}$  ( $0.0069 \mu\text{g insect}^{-1}$ ), i.e., 40.6 times more toxic than the lowest values recorded for malathion ( $0.28 \mu\text{g insect}^{-1}$ ). At the  $\text{LD}_{95}$ , however, pirimiphos-methyl ( $0.0018 \mu\text{g insect}^{-1}$ ) demonstrated the highest toxicity, being as much as 3,300 times more toxic than malathion at the lowest level ( $5.97 \mu\text{g insect}^{-1}$ ) of the tested insecticides.

**Table 1** Insecticide toxicity to *T. castaneum* adults from a laboratory population assayed by topical application after 7 d of exposure.

Insecticide	$\text{LD}_{50}$ ( $\mu\text{g insect}^{-1}$ ) Fiducial Limits (0.05)	$\text{LD}_{95}$ ( $\mu\text{g insect}^{-1}$ ) Fiducial Limits (0.05)	Slope <i>ld-p</i> line ( $\pm\text{SE}$ )	$\text{DD}^*$ ( $\mu\text{g insect}^{-1}$ )
Dichlorvos	0.040 (0.038-0.043)	0.087 (0.079-0.097)	4.98 $\pm$ 0.34	0.12
Malathion	0.28 (0.21-0.36)	5.97 (4.08-9.66)	1.24 $\pm$ 0.08	10.0
Chlorpyrifos-methyl	0.011 (0.0104-0.0115)	0.019 (0.018-0.021)	6.52 $\pm$ 0.42	0.030
Pirimiphos-methyl	0.0091 (0.0087-0.0095)	0.0018 (0.017-0.019)	5.68 $\pm$ 0.27	0.025
Deltamethrin	0.0069 (0.0062-0.0075)	0.024 (0.021-0.027)	3.05 $\pm$ 0.15	0.050
Bifenthrin	0.049 (0.041-0.056)	0.27 (0.22-0.35)	2.19 $\pm$ 0.16	0.50

\*Discriminating dose

Discriminating doses of the insecticides dichlorvos, pirimiphos-methyl, chlorpyrifos-methyl, deltamethrin and bifenthrin (Table 2) caused high mortality in all collected populations of *T. castaneum*, ranging from 92-100%. The discriminating dose of malathion caused 100% mortality only in a population from Bačka Topola (flour mill), and high mortality in another Bačka Topola population (silo) and one from Belgrade Port (silo), 95 and 99%, respectively. Mortality in all other populations was below 85%, and the lowest in the Jakovo population (silo), up to 64%.

**Table 2** Effects of discriminating doses of insecticides to *T. castaneum* adults from different populations assayed by topical application after 7 d of exposure.

Population	Mortality (%±SE)					
	DDVP	PM	CPM	MAL	DEL	BIF
Bačka Topola-FM	99±0.5	99±0.5	100	100	94±1.3	96±0.8
Bačka Topola-S	97±1.0	93±1.0	92±0.8	95±0.5	96±0.8	100
Belgrade Port-S	99±0.5	93±1.5	100	99±0.5	96±1.2	100
Čurug-S	99±0.5	92±2.2	99±0.5	80±0.8	98±0.6	100
Nikinci-W	98±0.6	94±1.3	100	70±2.5	99±0.5	100
Jakovo-S	100	99±0.5	100	64±2.9	100	100
Adaševci-S	99±0.5	97±0.5	100	79±2.2	100	100
Novo Miloševo-S	99±0.5	99±0.5	98±0.6	80±0.8	98±0.6	100
Irig-S	93±1.0	94±0.6	96±1.2	74±2.4	98±0.6	97±1.0
Kikinda-FM	100	99±0.5	98±1.0	84±2.2	96±1.4	100

S-silo; W-warehouse; FM-flour mill

Tables 3 and 4 show the results of insecticide toxicity (LD levels and *ld-p* lines) and resistance ratios (RRs) for beetles from Nikinci and Jakovo. For the *T. castaneum* from Nikinci, deltamethrin was the most toxic insecticide (0.0066 µg insect<sup>-1</sup>) at the LD<sub>50</sub>, and chlorpyrifos-methyl (0.016 µg insect<sup>-1</sup>) was most toxic at the LD<sub>95</sub>, which were 748.5 and 4097.5 times more toxic, respectively, than the lowest value for malathion (LD<sub>50</sub> 4.94 µg insect<sup>-1</sup> and LD<sub>95</sub> 65.56 µg insect<sup>-1</sup>).

For *T. castaneum* adults from Jakovo deltamethrin (0.0054 µg insect<sup>-1</sup>) was the most toxic insecticide at the LD<sub>50</sub>, while pirimiphos-methyl (0.0163 µg insect<sup>-1</sup>) was most toxic at the LD<sub>95</sub>, and they were 1348.1 and 2238.0 times more toxic, respectively, than the lowest toxicity for malathion (LD<sub>50</sub> 7.28 µg insect<sup>-1</sup> and LD<sub>95</sub> 36.48 µg insect<sup>-1</sup>). The RRs show that both populations were resistant to malathion as its RRs were 17.6 and 26.0 at the LD<sub>50</sub>, and 11.0 and 6.1 at the LD<sub>95</sub> for beetles from Nikinci and Jakovo, respectively.

**Table 3** Insecticide toxicity to *T. castaneum* adults from Nikinci population assayed by topical application after 7 d of exposure and resistance ratios found (RR = LD tested/LD laboratory population).

Insecticide	LD <sub>50</sub> (µg insect <sup>-1</sup> ) Fiducial Limits (0.05)	RR for LD <sub>50</sub> level	LD <sub>95</sub> (µg insect <sup>-1</sup> ) Fiducial Limits (0.05)	RR for LD <sub>95</sub> level	Slope <i>ld-p</i> line (±SE)
Dichlorvos	0.042 (0.039-0.044)	1.0	0.087 (0.080-0.096)	1.0	5.15±0.33
Malathion	4.94 (4.23-5.72)	17.6	65.56 (49.25-93.58)	11.0	1.46±0.01
Chlorpyrifos-methyl	0.0098 (0.0093-0.010)	0.9	0.016 (0.015-0.018)	0.8	7.21±0.60
Pirimiphos-methyl	0.011 (0.010-0.011)	1.2	0.025 (0.022-0.029)	1.4	4.48±0.34
Deltamethrin	0.0066 (0.0058-0.0073)	0.9	0.029 (0.025-0.035)	1.2	2.53±0.17
Bifenthrin	0.054 (0.049-0.060)	1.1	0.14 (0.11-0.17)	0.5	4.07±0.37

**Table 4** Insecticide toxicity to red flour beetle adults from Jakovo population assayed by topical application after 7 d of exposure and resistance ratios found (RR = LD tested/LD laboratory population).

Insecticide	LD <sub>50</sub> (µg insect <sup>-1</sup> ) Fiducial Limits (0.05)	RR for LD <sub>50</sub> level	LD <sub>95</sub> (µg insect <sup>-1</sup> ) Fiducial Limits (0.05)	RR for LD <sub>95</sub> level	Slope <i>ld-p</i> line (±SE)
Dichlorvos	0.035 (0.033-0.037)	0.8	0.079 (0.071-0.090)	0.9	4.65±0.34
Malathion	7.28 (6.54-8.03)	26.0	36.48 (30.85-44.85)	6.1	2.35±0.14
Chlorpyrifos-methyl	0.012 (0.0113-0.0127)	1.1	0.024 (0.022-0.027)	1.3	5.55±0.40
Pirimiphos-methyl	0.0090 (0.0085-0.0094)	0.9	0.0163 (0.0154-0.0176)	0.9	6.31±0.38
Deltamethrin	0.0054 (0.0047-0.0060)	0.7	0.024 (0.020-0.029)	1.0	2.55±0.19
Bifenthrin	0.040 (0.034-0.045)	0.8	0.17 (0.14-0.21)	0.6	2.65±0.23

#### 4. Discussion

As in many other studies worldwide, our experiments confirmed resistance to malathion. This is based on the effects of discriminating doses of contact insecticides on the ten tested populations of *T. castaneum*, and toxicity parameters and RRs for populations from Nikinci and Jakovo. However, compared to data on species resistance to insecticides summarized by Subramanyam and Hagstrum (1996) and Kljajić and Perić (2005), the level of resistance to malathion was low in the two examined populations in our experiment (RRs 17.6 and 26.0 at the LD<sub>50</sub> for beetles from Nikinci and Jakovo, respectively).

Subramanyam et al. (1989) had examined the susceptibility of *T. castaneum* and *Oryzaephilus surinamensis* (L.) (Coleoptera: Silvanidae) populations collected from barley warehouses using discriminating doses of malathion, pirimiphos-methyl and chlorpyrifos-methyl applied to filter paper. All tested populations of *T. castaneum* displayed resistance to malathion, but no cross resistance to either pirimiphos-methyl or chlorpyrifos-methyl.

In a study by Halliday et al. (1988) using topical application of discriminating doses of malathion, dichlorvos, pirimiphos-methyl, chlorpyrifos-methyl and synergized pyrethrins, several populations of *T. castaneum* from stored peanuts were resistant to malathion alone, mortality in 12 of the 15 tested population below 10%. Zettler (1991) used the same method to test susceptibility of *T. castaneum* and *Tribolium confusum* Jacquelin du Val populations collected from mills in the USA to malathion, dichlorvos, chlorpyrifos-methyl, synergized pyrethrins and resmethrin. Of the 28 tested populations of *T. castaneum*, 93% were resistant to malathion, 64% to dichlorvos, 36% to chlorpyrifos-methyl, while none were resistant to synergized pyrethrins or resmethrin.

More recently, Zettler and Arthur (1997) tested the susceptibility of 14 populations of *T. castaneum* and 10 populations of *T. confusum* from various mills to malathion and dichlorvos and found all *T. castaneum* populations to be resistant to dichlorvos and even more to malathion (top RR=29,081). They found a positive correlation between insect survival after exposure to discriminating doses of insecticides and lethal dose values.

In conclusion, our results confirmed the importance monitoring the susceptibility/resistance of *T. castaneum* and other stored-product insects to insecticides as an important element of pest management programs.

#### Acknowledgement

This study was supported by the Serbian Ministry of Science and Technological Development (Project number: TR-20060).

## References

- Abbott, W.S., 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18, 265-267.
- Almaši, R., 2008. Štetne artropode uskladištenog žita. U: Zaštita uskladištenih biljnih proizvoda od štetnih organizama (Kljajić, P., ed), Institut za pesticide i zaštitu životne sredine, Beograd, 9-38. (Harmful arthropods in stored grains and grain products, In: Kljajić, P. (Ed), Protection of Stored Plant Products from Harmful Organisms, Institute of Pesticides and Environmental Protection, Belgrade (in Serbian with abstract in English).
- Champ, B.R., Dyte, C.E., 1976. Report of the FAO Global survey of pesticide susceptibility of stored grain pests. FAO Plant Production and Protection Series No. 5, Food and Agriculture Organization (FAO), Rome, Italy, pp. 297.
- Davis, R., Bry, R.E., 1985. *Sitophilus granarius*, *Sitophilus oryzae* and *Sitophilus zeamais*; *Tribolium confusum* and *Tribolium castaneum*. In: Singh, P., Moore, R.F. (Eds), Handbook of Insect Rearing, Elsevier, Amsterdam, The Netherlands, pp. 287-293.
- Finney, D.J., 1971. Probit Analysis, Third edition Cambridge University Press, Cambridge, UK.
- Halliday, W.R., Arthur, F.H., Zettler, J.L., 1988. Resistance status of red flour beetle (Coleoptera: Tenebrionidae) infesting stored peanuts in southeastern United States. *Journal of Economic Entomology* 81, 74-77.
- Harein, C.R., Soderstrom, E.L., 1966. Coleoptera infesting stored products. In: Smith, C.N. (Ed), Insect colonization and mass production, Academic Press, New York, NY, USA, pp. 241-257.
- Kljajić, P., 2008. Suzbijanje štetnih insekata uskladištenog žita. U: Zaštita uskladištenih biljnih proizvoda od štetnih organizama (Kljajić, P., ed). Institut za pesticide i zaštitu životne sredine, Beograd, 101-122. (Control of harmful insects in stored grains. In: Kljajić, P. (Ed), Protection of Stored Plant Products from Harmful Organisms, Institute of Pesticides and Environmental Protection, Belgrade Serbia (in Serbian with abstract in English).
- Kljajić, P., Perić, I., 2005. Resistance of stored-product insects to insecticides (Rezistentnost skladišnih insekata prema insekticidima). *Pesticide & Phytomedicine (Pesticidi i fitomedicina)* 20, 9-28. (in Serbian with abstract in English).
- Raymond, M., 1985. Presentation d'un programme "Basic" d'analyse log-probit pour micro-ordinateur. Cahier ORSTOM, série. Entomologie, Médecine et Parasitologie 22, 117-121.
- Rees, D.P., 2004. Insects of Stored Products. CSIRO Publishing, Collingwood, Australia.
- Subramanyam, Bh., Hagstrum, D.W., 1996. Resistance measurement and management. In: Subramanyam, Bh., Hagstrum, D.W. (Eds), Integrated Management of Insects in Stored Products. Marcel Dekker Inc., New York, NY, USA, pp. 331-397.
- Subramanyam, Bh., Harein, P.K., Cutkomp, L.K., 1989. Organophosphate resistance in adults of red flour beetle (Coleoptera: Tenebrionidae) and sawtoothed grain beetle (Coleoptera: Cucujidae) infesting barley stored on farms in Minnesota. *Journal of Economic Entomology* 82, 989-995.
- Zettler, J.L., 1991. Pesticide resistance in *Tribolium castaneum* and *T. confusum* (Coleoptera: Tenebrionidae) from flour mills in the United States. *Journal of Economic Entomology* 84, 763-767.
- Zettler, J.L., Arthur, F.H., 1997. Dose-response tests on red flour beetle and confused flour beetle (Coleoptera: Tenebrionidae) collected from flour mills in the United States. *Journal of Economic Entomology* 90, 1157-1162.
- Zettler, J.L., Cuperus, G.W., 1990. Pesticide resistance in *Tribolium castaneum* (Coleoptera: Tenebrionidae) and *Rhyzoperha dominica* (Coleoptera: Bostrichidae) in wheat. *Journal of Economic Entomology* 83, 1677-168.