

Determination of fipronil LD₅₀ for the brazilian bee *Melipona scutellaris*

Clara T. Lourenço¹, Stephan M. Carvalho², Osmar Malaspina², Roberta C. F. Nocelli^{1*}

¹CCA - Centro de Ciências Agrárias, UFSCar - Universidade Federal de São Carlos . Rod. Anhanguera, SP 330, Km. 174, Araras – SP, Brasil. Email: claratavareslourenco@yahoo.com.br; Email: roberta@cca.ufscar.br,

*Phone: +55 (19) 3543-2595

²CEIS – Centro de Estudos de Insetos Sociais. Universidade Estadual “Júlio de Mesquita Filho” - UNESP. Av. 24 A, 1515, Bela Vista, Rio Claro – SP, Brasil.

DOI: 10.5073/jka.2012.437.046

Abstract

To better understand the sensitivity of the models represented by *Apis mellifera* L., 1758 in toxicology studies of insecticides to bees, the aim of this study was to determine the LD₅₀ of fipronil by topical application on the stingless bees *Melipona scutellaris* Latreille, 1811. Foraging bees were collected at the nest entrance and in laboratory anesthetized with CO₂ for applying 1.0 µL of fipronil solution on the pronotum. Each group of treatments was made with thirteen bees divided in three cages, while in the control treatments the bees received only acetone. During the assay, the behavior and the number of dead bees were registered. The results showed that the insecticide fipronil applied topically was harmful to *M. scutellaris* and for *A. mellifera*, where the LD₅₀ for 48 hours was 0.41 ng a.i./bee or 4.1 ng a.i./ g of bee. Comparing the LD₅₀ values here obtained with the stingless bee *M. scutellaris* and those of *A. mellifera* in literature, we can conclude that the native bees are more sensitive to fipronil than the allochthonous bee, suggesting that further studies should be accomplished to determine the real hazard of pesticides to natives bees.

Keywords: stingless bees, pesticides, phenylpyrazole, toxicity, LD₅₀

1. Introduction

The insecticide fipronil (phenylpyrazole - C₁₂H₄Cl₂F₆N₄O₅) acts on the nervous system of insects by blocking chloride channels through the receptor of gamma-aminobutyric acid and glutamate. Due to its mode of action it is considered a new generation insecticide, being a broad-spectrum systemic insecticide and effective at low application rates. Widely used in Brazil and more than 70 countries it is considered highly toxic to bees, which is why its use is banned in France since 2004¹⁻¹⁰.

Toxicological studies with bees use mostly the model species *Apis mellifera* L., 1758 (Hymenoptera: Apidae) and it has been observed that sublethal doses of fipronil can cause behavioral changes, to these bees, related mainly with tasks such as feeding and foraging, fundamental for the survival of colonies¹¹⁻¹⁸. However, native bees exposed to pesticides may be at a different risk, because differences in tolerance among species of bees have been observed by several authors¹⁹⁻²⁶ and most show that wild bees are more sensitive to the insecticides than the honeybee *A. mellifera*^{27,28}.

Among the stingless bees within the tribe Meliponini, *Melipona scutellaris* Latreille, 1811 (Hymenoptera: Apidae) is popularly known. It is endemic in northeastern Brazil and distinguished by its ease of domestication and management, honey production and significant potential for replication on a large scale for pollination in greenhouses and open field, beyond its ecological importance as pollinators of native plants in Brazil²⁹⁻³⁴.

Seeking to diversify the species of bees to better understand and compare the sensitivity of the models represented by honeybee *A. mellifera*²⁷, the objective of this study was to determine the topical LD₅₀ of the insecticide fipronil for the stingless bees *M. scutellaris*.

2. Material and Methods

Three colonies of *M. scutellaris* from the Universidade Estadual Paulista (UNESP) campus Rio Claro, were used in the experiment. The hives were kept in protected room where the bees had free access to the external environment through a plastic tube that connected the nest entrance and the outside. All time, the colonies were surveyed to assess the health, queen laying capacity, foraging activity and

food availability. To promote the survival of colonies during a dry season, 60 % of sucrose solution prepared with lemon juice was provided³⁵.

2.1 Acute toxicity test (LD₅₀)

Assays were carried out at the Center for the Study of Social Insects of UNESP with some modifications on the directives of the Organization for Economic Cooperation and Development³⁶.

To determine the topical LD₅₀ of the insecticide fipronil (95% of purity, Bayer CropScience, Brazil) to foragers of *M. scutellaris*, a stock solution (1000 ng a.i./μL acetone) was prepared and next a range of several concentrations between 0.5 to 5.0 ng of a.i./μL acetone. The control treatment received only acetone, after its low toxicity had been assessed in preliminary assay compared with water.

To facilitate the handling and application, the bees were anesthetized with CO₂ (ten seconds). With a repetitive micropipette the volume of 1.0 μL of solution was applied on the thorax of the bees. To ensure the variability among the colonies and to obtain a realistic and reliable value of LD₅₀, the bees from each repetition were taken directly from a single colony. In each treatment (=concentration) we had three distinct groups of bees which originated from different colonies. Thus, each treatment (group of bees from one concentration) consisted of three replicates with ten bees, in total thirty specimens. During the assay, bees were fed *ad libitum* with sucrose-solution (50%), and cages were kept in climatic room at 29±2°C, relative humidity of 70±5% and darkness.

2.2 Data collection and analysis

Along 72 hours after the application of fipronil on *M. scutellaris*, assessments were made one, four and every twenty-four hours, with registration of all behavior, as well as the number of dead bees. Statistical analysis to determine the LD₅₀ value were performed using a log-logistic model from the package "drc" (Analysis of Dose-Response Curves)³⁹ compiled by the statistical software R⁴⁰.

3. Results and discussion

Bioassays performed in order to compare the toxicity of acetone and water showed that this organic solvent was no toxic to *M. scutellaris* foragers. Already, the insecticide fipronil topically applied was considered highly toxic to *M. scutellaris* foragers, with a LD₅₀ for 48 hours of 0.41 ng a.i./bee (CL_{95%} = 0.23 - 0.58; D.F. = 16 and $\chi^2 = 9.8238$, Figure 1). Comparing this result with the LD₅₀ of fipronil established for other species of bees, foragers of *M. scutellaris* were more sensitive to fipronil than *A. mellifera* (1.9- 6 ng a.i./bee), *Megachile rotundata* Fabricius, 1787 (4 ng a.i./bee), *Nomia melanderi* Cockerell, 1906 (113 ng a.i./bee) and *Scaptotrigona postica* Latreille, 1807 (0.54 ng a.i./bee)^{13,18,37,38,41, 42}. Likewise, taking into consideration that the workers of *M. scutellaris* have a mean weight of 0.1g, recalculating the LD₅₀ we got a LD₅₀ of 4.1 ng a.i./g of bee. In this way, *M. scutellaris* remains more susceptible to fipronil than *A. mellifera* (103 ng a.i./g of bee), *M. rotundata* (132 ng a.i./g of bee) and *N. melanderi* (13,190 ng a.i./g of bee)⁴¹.

The doses of fipronil from 1.5 to 5 ng a.i./bee for 48/72 hours and 5 ng a.i./bee for 24 hours, caused 100 % of mortality, respectively (Figure 2) The dose of 1.5 and 1.0 ng / bee also had high rates of mortality after 48 hours of intoxication, with 96% and 85% of dead bees, respectively.

Still, the bees in the group treated with 5.0 ng of fipronil/bee showed signs of intoxication: after 4 hours this group had bees with their wings vibrating. This same behavior was observed in *A. mellifera* treated with 0.1 ng fipronil/bee and after 11 days of exposure dose of 0.01 ng of fipronil/bee¹¹. According to these authors the behavior of vibrating wings is accompanied by the emission of alarm pheromone, which causes attacks among individuals, also observed in this study. After 24 hours of contamination, surviving bees in the groups treated with the higher doses of fipronil (2.0, 2.5 and 5.0 ng a.i./bee) had tremors followed by paralysis and death. These same signals were also observed in honeybee *A. mellifera* treated with sucrose solution contaminated with fipronil 2 g/Kg of diet¹².

These results are consistent among the diversity of bees, which differ in their vulnerability to exposure to insecticides²⁸. Several studies¹⁹⁻²⁶ show differences in tolerance and/or sensitivity between species of bees and pesticides, most of these results show that the species honeybee *A.*

mellifera was more resistant compared to species of stingless bees which corroborates the suggestion that wild bees are a pollinating group at particular risk for exposure to pesticides²⁷.




Fig. 1 Acute toxicity (48 hours) by topical application of the insecticide fipronil to foragers of *Melipona scutellaris*.




Fig. 2 Mortality evolution of foragers *Melipona scutellaris* when exposed at different doses of fipronil.

The findings also support the idea that ecotoxicological studies on diverse pollinating species can be used to obtain a better understanding of how the sensitivity of a model representative of honeybees (*A. mellifera*) can be compared to other species of bees⁴³.

The bee species *M. scutellaris* is considered a potential effective pollinator species for production on a large scale as a pollinator in greenhouses and the open field, with ease of maintaining strong hives, which can be easily transported and multiplied³⁰. Brazil has a high diversity of bees that interact with numerous plant species⁴⁴ and it is believed that 33% of the crops that provide food for the human population depends on pollination by bees⁴⁵.

We believed than poisoning by insecticides is one of the causes of high mortalities of bees, especially in areas in southern and southeastern Brazil, where the disappearance of bees caused by insecticides

has become a concern. Between 2008 and 2010, about 5000 bee hives of Africanized *A. mellifera* were lost in the central region of São Paulo. Hives of native bees were not included⁴⁶.

Since Brazil has a high diversity of native bees, endemic in the tropics and sensitive to low temperatures, studies on the toxicity of insecticides in Brazil should focus on these species⁴⁷.

4. Conclusion

The insecticide fipronil was highly toxic to foraging stingless bee *M. scutellaris* under laboratory conditions, with a topical LD₅₀ in 48 hours of 0.41 ng a.i./bee (4.1 ng of fipronil/g bee). It is suggested that bees of *M. scutellaris* are more sensitive to fipronil than *A. mellifera* (Africanized and Italian), *M. rotundata*, *N. melanderi* and *S. postica*. Tremor followed by paralysis were the main signs of intoxication observed in the groups treated topically with the highest dose of fipronil.

The LD₅₀ results determined in this work are being used to assess behavioral changes through the Proboscis Extension Reflex (PER) and locomotor activity, in particular which doses of fipronil causing sublethal effects in foragers of *M. scutellaris*.

Acknowledgement

This study was supported by CAPES (Brazil), Dutch Ministry of Agriculture, Nature and Food Quality (BO-10-011-113) and GEF.

References

1. Aajoud A, Ravanel P, Tissut M, Fipronil metabolism and dissipation in a simplified aquatic ecosystem, *J Agric Food Chem* **51**: 1347-1352 (2003).
2. Agência nacional de vigilância sanitária/Anvisa, fipronil, [http://www4.anvisa.gov.br/base/visadoc/cp/cp\[18765-1-0\].pdf](http://www4.anvisa.gov.br/base/visadoc/cp/cp[18765-1-0].pdf) [accessed on September 2011].
3. Barbara GS, Zube C, Rybak J, Gauthier M, Grünewald B, Acetylcholine, Gaba and glutamate induce ionic currents in cultured antennal lobe neurons of the honeybee, *Apis mellifera*, *J Comp Physiol A*, **191**: 823-836 (2005).
4. Connely P, Environmental fate of fipronil. Environmental monitoring branch, department of pesticide regulation, California Environmental Protection Agency, <http://www.pw.ucr.edu/textfiles/fipronil.pdf> [accessed on September 2011].
5. Gunasekara AS, Truong T, Goh KS, Spurlock F, Tjeerdema RS, Environmental fate and toxicology of fipronil, *J Pestic Sci*, **32**(3): 189-199 (2007).
6. Johnson R, Honey bee colony collapse disorder, *Congressional Research Service*, 20 p. (2010).
7. National Pesticide Information Center/Npic, Fipronil: technical fact sheet, <http://npic.orst.edu/factsheets/fiptech.pdf> [accessed on January 2010].
8. Rogers REL, Why is France banning certain pesticides?, *Bee World*, p. 40-41, jun. 2004.
9. Stenersen J, Interference with signal transduction in the nerves. In: Chemical Pesticides: Mode of action and toxicology. Crc Press, New York, (2004).
10. Thompson HM, Risk assessment for honey bees and pesticides – recent development and 'new issues', *Pest Manag Sci*, **66**: 1157-1152 (2010).
11. Aliouane Y, Hassani AK, Gary V, Armengaud C, Lambin M, Gauthier M, Subchronic exposure of honeybees to sublethal doses of pesticides: effects on behavior, *Environ Toxicol Chem*, **28** (1): 113-122 (2009).
12. Colin ME, Bonmatin JM, Moineau I, Gaimon C, Brun S, Vermandere JP, A method to quantify and analyse the foraging activity of honey bees: relevance to the sublethal effects induced by systemic insecticides, *Arch Environ Contam Toxicol*, **47**: 387-395 (2004).
13. Decourtye A, Devillers J, Genecque E, Menach K, Budzinski H, Cluzeau S, Pham-Dèlegue MH, Comparative sublethal toxicity of nine pesticides on olfactory learning performances of the honeybee *Apis mellifera*. *Arch Environ Contam Toxicol*, **48**: 242-250 (2005).
14. Decourtye A, Lefort S, Devillers J, Gauthier M, Aupinel P, Tisseur M, Sublethal effects of fipronil on the ability of honeybees (*Apis mellifera* L.) to orientate in a complex maze. *Julius Kühn Arch*, **423**: 75-83 (2009).
15. Decourtye A, Devillers J, Aupinel P, Brun F, Bagnis C, Fourrier J, Gauthier M, Honeybee tracking with microchips: a new methodology to measure the effects of pesticides, *Ecotoxicol*, **20**: 429-437 (2011).
16. Hassani AK, Dacher M, Gauthier M, Armengaud C, Effects of sublethal doses of fipronil on the behavior of the honeybee (*Apis mellifera*), *Pharmacol Biochem Behavior*, **82**(1): 30-39 (2005).

17. Hassani AK, Dupuis JP, Gauthier M, Armengaud C, Glutamatergic and gabaergic effects of fipronil on olfactory learning and memory in the honeybee, *Invert Neurosci*, **9**(2): 91-100 (2009).
18. Pereira AM, unpublished.
19. Del Sarto MCL, unpublished.
20. Macieira OJD, Hebling-Beraldo MJA, Laboratory toxicity of insecticides to workers of *Trigona spinipes* (F., 1793) (Hymenoptera, Apidae). *J. Apic Res*, **28**(1): 3-6 (1989).
21. Malaspina O, pers. communication.
22. Malaspina O, Stort AC, DDT tolerance of africanized bees, italian bees (*Apis mellifera ligustica*) and their F1 hybrids (Hymenoptera: Apidae), *J. Kansas Entomol Soc*, **56**(1): 74-79 (1983).
23. Moraes SS, Bautista ARL, Viana BF, Avaliação da toxicidade aguda (DL₅₀ E CL₅₀) de inseticidas para *Scaptotrigona tutiba* (Smith) (Hymenoptera: Apidae): via de contato, *An Soc Entomol Brasil*, **29**(1): 31-37 (2000).
24. Roessink I, Steen J, Kasina M, Gikungu M, Nocelli R, Is the european honeybee (*Apis mellifera mellifera*) a good representative for other pollinators species?, *Setac*, p. 35 (2011).
25. Stuchi ALPB, unpublished.
26. Valdovinos-Núñez GR, Quezada-Euán JGG, Ancona-Xiu P, Moo-Vale H, Carmona A, Sánchez ER, Comparative toxicity of pesticides to stingless bees (Hymenoptera: Apidae: Meliponini), *J Econ Entomol*, **102**(5) 1737-1742 (2009).
27. Brittain CA, Vighi M, Bommarco R, Sattelle J, Potts SG, Impacts of a pesticide on pollinators species richness at different spatial scale, *Basic Appl Ecol*, **11**(2): 106-115 (2010).
28. Desneux N, Decourtye A, Delpuech JM, The sublethal effects of pesticides on beneficial arthropods, *Annu Rev Entomol*, **52**: 81-106 (2007).
29. Evangelista-Rodrigues A, Góis GC, Silva CM, Souza DL, Souza DN, Silva PCC, Alves EL, Rodrigues ML, Desenvolvimento produtivo de colméias de abelhas *Melipona scutellaris*, *Biotemas*, **21**(1): 59-64 (2008).
30. Imperatriz-Fonseca VL, Saraiva AM, De Jong D, Bees as pollinators in Brazil: assessing the status and suggesting best practices. Holos, Ribeirão Preto, 112p. (2006).
31. Kerr WE, As abelhas e o meio ambiente. 12 Congresso Brasileiro de Apicultura, Anais, p. 1-8 (1999).
32. Kerr WE, Carvalho GA, Silva AC, Assis MGP, Aspectos pouco mencionados da biodiversidade Amazônica, *Parcerias Estrateg*, **12**: 20-41 (2001).
33. Nogueira-Neto P, Vida e criação de abelhas indígenas sem ferrão. Nogueirapis, São Paulo, 445 p. (1997).
34. Silveira FA, Melo GAR, Almeida EAB, Abelhas Brasileiras: sistemática e identificação. Fernando a. Silveira, Belo Horizonte, 253 p. (2002).
35. Brighenti DM, Carvalho CF, Brighenti CRM, Carvalho SM, Inversão da sacarose utilizando ácido cítrico e suco de limão para preparo de dieta energética de *Apis mellifera* Linnaeus, 1758, *Cienc Agrotec*, **35**(2): 297-304 (2011).
36. Organization for Economic Cooperation and Development/OECD. Honeybees, acute contact toxicity test, OECD guidelines for the testing of chemicals, 8 p. (1998).
37. Roat TC, pers. communication.
38. Jacob CRO, pers. communication.
39. Ritz C, Streibig JC, Bioassay analysis using *r*, *J Statistical Software*, **12**(5): 1-22 (2005).
40. R development core team. R: A language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria (2011). ISBN 3-90051-07-0, URL <http://www.R-project.org> [accessed on October 2011].
41. Mayer DF, Lunden JD, Field and laboratory tests of the effects of fipronil on adults female bees of *Apis mellifera*, *Megachile rotundata* and *Nomia malarandi*, *J Apic Res*, **38**(3-4): 191-197 (1999).
42. Tingle CC, Rother JA, Dewhurst CF, Lauer S, King WJ, Fipronil environmental fate, ecotoxicology and human health concerns, *Rev Environ Contam Toxicol*, **176**:1-66 (2003).
43. Brittain C, Potts SG, The potential impacts of insecticides on the life-history traits of bees and the consequences for pollination, *Basic Appl Ecol*, **12**: 321-331 (2011).
44. Klein AM, Vaissière BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C, Tscharntke T, Importance of pollinators in changing landscapes for world crops, *Proc R Soc B*, **274**: 303-313 (2007).
45. Imperatriz-Fonseca VL, Silva-Nunes P, As abelhas, os serviços ecossistêmicos e o código florestal brasileiro, *Biota Neotrop*, **10**(4): 59-62 (2010).
46. Malaspina O, Nocelli RCF, Silva-Zacarin ECM, Souza TF, Defesa de apiários e meliponários contra agrotóxicos, 18 Congresso Brasileiro de Apicultura, 4 Congresso Brasileiro de Meliponicultura, Anais, 5p. (2010).
47. Lourenço CT, Santos JF, Nocelli RCF, Effects of insecticides in bees: situation in Brazil, 9 Encontro sobre abelhas, ed by FFCLRP-USP, Ribeirão Preto, p. 607 (2010).