

## Reduction of in-shell Brazil nut (*Bertholletia excelsa* H.B.K.) aflatoxin contamination by ozone gas application during storage

Giordano, B.N.E.<sup>1</sup>, Simao, V.<sup>1</sup>, Manfio, D.<sup>1</sup>, Galvao, S.<sup>1</sup>, Scussel, J.N.<sup>2</sup>, Scussel, V.M.\*<sup>#1</sup>

<sup>1</sup> Food Science & Technology Department, Center of Agricultural Sciences, Federal University of Santa Catarina, P.O.Box 476, Florianopolis, SC, CEP 88034-001, Brazil. Email: vildescussel\_2000@yahoo.co.uk

<sup>2</sup> Mechanical Engineering Dept Technological Center, Federal University of Santa Catarina, PO.Box 476, Florianopolis, SC, Brazil.

\* Corresponding author

# Presenting author

DOI: 10.5073/jka.2010.425.167.231

### Abstract

The susceptibility of the in-shell Brazil nut mycoflora and aflatoxins (AFLs) contamination to ozone (O<sub>3</sub>) gas during storage is reported. In-shell Brazil nuts obtained from retail market were submitted to O<sub>3</sub> gas atmosphere at different concentrations immediately before to be stored. Samples were collected just after the gas exposure and every 30 days during the storage period to carry on mycological tests and AFLs analysis. A sensorial evaluation by descriptive quality analysis was carried out to check treated nuts sensory attributes according to consumer acceptance after gas exposure. The O<sub>3</sub> treatment applied within 5 h at 31 mg/L was able to successfully inhibit the viability of fungi of the nut-contaminating microflora and so the toxigenic *Aspergillus species* from the day of application. AFLs were totally degraded in all samples whatever O<sub>3</sub> concentration applied. No significant changes on sensory attributes were observed that could affect nut acceptability after the O<sub>3</sub> treatments and storage conditions applied in the present experiment. This procedure is tentatively applied at an Amazon State nut factory for checking its potential in mycotoxin risk contamination of in-shell Brazil nuts safeguarding under the Amazon region environment.

Keywords: In-shell Brazil nut, Ozone, Mycoflora, Aflatoxin, Storage, Sensory evaluation.

### 1. Introduction

Prevention of aflatoxin (AFL) contamination in tree nuts by controlling the environment conditions for toxigenic fungi growth has not always been effective. The risk of AFL contamination in Brazil nuts (*Bertholletia excelsa* H.B.K.) can occur, either in the forest or during storage and it can vary with the year of harvest (Pacheco and Scussel, 2006; 2009). Nut contamination by fungi, especially the aflatoxigenic species of *Aspergillus*, are directly related to the climatic conditions (high humidity and warm conditions) in the Amazon forest during the period of harvesting (at wet season: December to May) (Arrus et al., 2005; Pacheco and Scussel, 2006; 2007; 2009; Olsen et al, 2008). The presence of AFLs is a concern for exporters of Brazil nuts according to European Community (EC) reduction the maximum tolerance limit of aflatoxins (AFB<sub>1</sub> + AFB<sub>2</sub> + AFG<sub>1</sub> + AFG<sub>2</sub>) and AFB<sub>1</sub> to 4 and 2 µg/kg, respectively (ECC, 2006) either in-shell or shelled. Thus, there is a need for the development of safe technologies at the forest environment level enabling to control fungi proliferation and reduce toxin contamination.

Fungi can growth both, on the shell and inside the shell on the nuts when shells are cracked or through its opercule. Thus, AFLs have been detected on the surface of shelled nuts and/or inside of cracked, or brown spotted, in-shell nuts (De Mello and Scussel, 2007; 2009). Recently, it was found that the main site of the in-shell nut contaminated by AFLs is located between the shell and the the nut peel (Conforcast, 2008). Several environmental factors are known to influence fungi growth and AFL production, being temperature and relative humidity (r.h.) the most critical (Mangan and Aldred, 2007). Ozonation, an oxidation method has been studied for detoxification of AFLs in foods (Shamarajeewa et al., 1990). The oxidation decreases AFL concentration over time (McKenzie et al., 1997). O<sub>3</sub> modified atmospheres have been developed for dried figs, barley, pistachio among other foods (Oztekin et al., 2006). The attractive aspect of O<sub>3</sub> is that it decomposes rapidly (half-life of 20–50 min) to molecular oxygen after application without leaving a residue (Mason et al., 1997; Frazier and Westhoff, 1988; Maeba et al., 1988; Perez et al., 1999; Kells et al., 2001; Yesilcimen and Murat, 2006; Inan et al., 2007; Olmez et al., 2009). It can be applied to food as a gas or dissolved in water.

One of the most important applications of O<sub>3</sub> in agriculture is in the post-harvest treatment of crops and the main purposes of gas application are: inactivation of bacterial growth, prevention of fungal decay, destruction of pesticides and chemical residues, and control of storage pests. Storage period can be doubled when some fruits and vegetables are held in an environment with O<sub>3</sub> (Frazier and Westhoff, 1988; Inan et al., 2007; Olmez et al., 2009). Regarding *fungi* and AFLs in different food commodities, Oztekin et al. (2006) reported a significant reduction of microorganism counts on dried figs at 5 mg/L of O<sub>3</sub>, decreasing the total yeast/mould counts of 72% and Perez, et al. (1999) observed a fungal decay of strawberry after 4 days of storage under ozonation. Five mg/L of O<sub>3</sub> inhibited the surface growth, sporulation and mycotoxin production of cultures of *Aspergillus flavus* and *Fusarium moniliforme* (Mason et al., 1997). Yesilcimen and Murat (2006) studied pistachio exposure to O<sub>3</sub>: 5.0, 7.0 and 9.0 mg/L and found AFB<sub>1</sub> and total AFLs reduction by 23 and 24% at 9 mg/L. Maeba et al. (1988) observed the destruction and detoxification of AFB<sub>1</sub> and AFG<sub>1</sub> with O<sub>3</sub> in model experiments.

We report here the application of O<sub>3</sub> gas at different times and concentrations during the storage of in-shell Brazil nut to improve nuts safety regarding fungi and AFLs contamination.


## 2. Materials and methods

### 2.1. Sample

Dry (processed) in-shell Brazil nuts (14 kg), export Medium Size Type 40-50 mm length (De Melo and Scussel, 2007). A naturally contaminated batch was chosen for the study with initial AFLs (AFB<sub>1</sub> / AFB<sub>2</sub> / AFG<sub>1</sub> / AFG<sub>2</sub>) of 11.58 µg/kg obtained in the retail market. That level is allowed by regulations from USA, Canada, Brasil and Mecosur (15, 15, 30, 20 µg kg, respectively). Total fungi count: 4.83 log cfu/g and 9.43% and m.c.

### 2.2. Storage

Seven vertical silos, build with vinyl polychloride (PVC) with the following dimensions 80 x 15 x 0.2 cm for height, diameter and width, respectively; containing an upper lid and two apertures (top and lower part of the silos) for sample collection and O<sub>3</sub> application, respectively; connected to an ozone generator (Megazon™) (Fig. 1).



**Figure 1** Brazil nut ozone treatment and storage system: silos (n=7) and O<sub>3</sub> generator

### 2.3. Chemicals, reagents and culture media

The reagents, potassium iodine, sulphuric acid, sodium thiosulfate, 2-thiobarbituric acid (TBA), trichloroacetic and butylated hydroxytoluene were of Analytical grade (Vetec) and utilized for the TBA test. The solvents were methanol, ethanol, acetonitrile, benzene and toluene from Carlo Erba, starch indicator from Synth and the ultrapure water from MilliQ system, Millipore. The standards of AFL

(AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub>) were obtained from Sigma. Malt extract agar-MEA was purchased from Himedia™; *A. flavus* and *parasiticus* agar-AFPA from Fluka™, peptone agar from Himedia™ and Tween 80 from CRQ™.

#### 2.4. Equipment and apparatus

The equipment and apparatus utilized for sample preparation, microbiological and AFL analysis were homogenizer from IKA T 25-Ultra Turrax; water bath from Quimis-Dubnoff™ Q226D; autoclave from Phoenix™; microscope from 100-400x PZO; incubator set at 20-25°C from Dist™; microscope stereoscope from Carl Zeiss Iena™ (Germany); colonies counter from Phoenix™ and microbiological oven from Fanen™. Analytical scales from Mettler™ and the semi-analytical one from CAB™. Industrial Brazil nut cracker was provided by CIEX™ (Manaus, AM, Brazil). The spectrophotometer model E005 was from Hitachi™. The liquid chromatograph comprised an isocratic pump model 305 and fluorescence detector model 121 from Gilson™. The column was of C<sub>18</sub> with 15 cm length, 4.6 mm diameter, and particle size of 5 µm from Phenomenex™. Thermometer and hygrometer utilized for environment temperature and moist readings were from CE™.

#### 2.5. Sample preparation for O<sub>3</sub> application

In-shell Brazil nuts previously analysed for fungi load, m.c. and AFLs contamination were aseptically weighted into portions of 2 kg to be added into each silo for the O<sub>3</sub> treatment.

#### 2.6. Preparation of the storage Silos, ozone application and iodometry

Silos were filled with nuts and tightly closed. They were divided into 4 groups for O<sub>3</sub> application: Group C (Control = no O<sub>3</sub> application), Group I (O<sub>3</sub> conc. = 10 mg/L), Group II (O<sub>3</sub> conc. = 14 mg/L) and Group III (O<sub>3</sub> conc. = 31.5 mg/L) (Fig. 1). Each treatment was carried out in duplicate (n = 2) except for Group C. After closing the upper part of the silos, O<sub>3</sub> gas was applied through the lower lateral aperture to get the following concentration in each silo: 10, 14 e 31.5 mg/L, with 1, 3 and 5 h exposure time, (n = 2) for silo Groups 2, 3 and 4, respectively. Those concentrations were checked by iodimetric analysis (APHA, 1980). The storage time was expanded on a 180-d period (May to October).

#### 2.7. Sample collection for analysis

Individual 200 g samples of Brazil nuts were aseptically collected from each silo at day-one and every 30 d. This sampling was carried out from the top silo aperture. Samples collected for analysis were made in duplicate (n = 2).

#### 2.8. Mycology, m.c., r.h., temperature, AFLs and sensory analysis

For total fungi count the method used was from Pitt and Hocking (1997). For fungi toxigenicity (for AFLs) the method was from Machida and Saito (1999). The identification of fungi in genus and species was carried out according to the keys of Samsom et al. (2004). The strains aflatoxigenicity was checked utilizing the AFPA method developed by Pitt et al. (1983). The m.c. was carried out by a gravimetric method (AOAC, 2005). Relative humidity and temperature were monitored daily utilizing hygrometer and thermometer, respectively. In parallel, data on r.h. and temperature was obtained from the National climatic recordings database EPAGRI/SC (May to October). AFLs analysis was performed by liquid chromatography with fluorescence-detection, with a limit of quantification (LOQ) at 0.50, 0.17, 0.50, and 0.17 µg/kg for AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub>, respectively (Sobolev et al., 2007).

#### 2.9. Sensory evaluation

The descriptive quantitative analysis by Stone and Sidel (1993) was conducted using a team of 18 trained (specifically for Brazil nut taste) panelists and four sessions (n = 4). The hedonic scale was separated into 5 degrees as follows: 1: dislike very much, 2: dislike, 3: neither like nor dislike, 4: like, and 5: like very much. Six sensory attributes of the Brazil nuts were recorded: shell appearance, nut appearance, strange odor, roasted flavor, rancidity and firmness.

### 2.10. Statistical analysis

The results were expressed as the mean values and standard errors. Statistical analysis was performed by analysis of variance (ANOVA) and included the Turkey's test to evaluate significant differences among the means ( $p < 0.05$ ).

## 3 Results and discussion

### 3.1. Effect of O<sub>3</sub> on the in-shell Brazil nut mycoflora

The total fungi load, aflatoxigenic *Aspergillus* species and levels of AFLs variation during the storage of in-shell Brazil nuts under O<sub>3</sub> atmosphere are shown in Table 1. There was a clear reduction on the total fungi count, AFLs and m.c. when compared to the Control Group. However this reduction trend was dependent upon the O<sub>3</sub> concentration used and time of storage.

**Table 1** Fungi, aflatoxigenic *Aspergillus* species and aflatoxins levels in Brazil nut stored under ozone at room temperature

Storage		Brazil nut											
days	O <sub>3</sub> treatment	Fungi				m.c./loss (%)		Aflatoxins (in-shell / after shelling) (µg/kg) <sup>a</sup>					
		total count (log cfu/g)		Aspergillus aflatoxigenic species		in-shell	after shelling	ΣAFLs	AFB <sub>1</sub>	AFG <sub>1</sub>	AFB <sub>2</sub>	AFG <sub>2</sub>	
		group	conc. (mg/L)	in-shell	after shelling								in-shell
1	C	0 <sup>b</sup>	4.83	2.54	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.43 (NA)	5.14 (NA)	11.58/6.61	3.48/1.16	3.57/1.89	2.21/2.02	2.32/1.74
	I	10	3.5	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.72 (6.61)	3.97 (6.25)	3.01/ND	3.01	ND	ND	ND
	II	14	3.3	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.71 (7.58)	3.96 (6.89)	ND	ND	ND	ND	ND
30	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.68 (8.56)	3.95 (7.52)	ND	ND	ND	ND	ND
	C	0	4.84	2.57	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.57 (NA)	5.28 (NA)	12.06/8.01	3.69/1.37	3.78/2.73	2.23/2.05	2.36/1.86
	I	10	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.67 (10.22)	3.95 (8.04)	ND	ND	ND	ND	ND
60	II	14	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.66 (10.60)	3.94 (8.67)	ND	ND	ND	ND	ND
	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.64 (11.74)	3.93 (9.29)	ND	ND	ND	ND	ND
	C	0	4.86	2.60	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.63 (NA)	5.32 (NA)	12.24/7.95	3.83/1.16	3.82/2.83	2.22/2.08	2.37/1.86
90	I	10	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.63 (11.65)	3.93 (9.03)	ND	ND	ND	ND	ND
	II	14	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.61 (12.59)	3.90 (9.65)	ND	ND	ND	ND	ND
	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.58 (13.34)	3.88 (10.28)	ND	ND	ND	ND	ND
120	C	0	4.88	2.62	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.84 (NA)	5.46 (NA)	12.34/8.03	3.86/1.14	3.86/2.86	2.25/2.10	2.37/1.86
	I	10	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.56 (14.65)	3.87 (11.38)	ND	ND	ND	ND	ND
	II	14	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.54 (15.75)	3.87 (12.29)	ND	ND	ND	ND	ND
180	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.51 (16.11)	3.60 (12.50)	ND	ND	ND	ND	ND
	C	0	4.89	2.65	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.89 (NA)	5.51 (NA)	12.49/8.08	3.94/1.14	3.88/2.88	2.27/2.11	2.40/1.86
	I	10	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.47 (16.15)	3.85 (12.23)	ND	ND	ND	ND	ND
180	II	14	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.43 (17.78)	3.84 (13.14)	ND	ND	ND	ND	ND
	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.41 (17.96)	3.82 (13.34)	ND	ND	ND	ND	ND
	C	0	4.91	2.69	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	9.93 (NA)	5.63 (NA)	12.55/8.17	3.95/1.13	3.90/2.91	2.28/2.17	2.42/1.86
180	I	10	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.37 (18.82)	3.80 (12.99)	ND	ND	ND	ND	ND
	II	14	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.35 (19.89)	3.78 (13.79)	ND	ND	ND	ND	ND
	III	31.5	ng	ng	<i>A. flavus</i> , <i>A. parasiticus</i>	ng ng	7.32 (19.89)	3.76 (14.19)	ND	ND	ND	ND	ND

<sup>a</sup> nuts were evaluated for fungi whole (in-shell + edible part) and after shelling (only edible part); nuts Groups: <sup>b</sup> C = control (no O<sub>3</sub> treatment); I (O<sub>3</sub> conc. = 10 mg/L), II (O<sub>3</sub> conc. = 14 mg/L) and III (O<sub>3</sub> conc. = 31.5 mg/L); NT = not treatment; NA = not applicable; ng = no growth; mc = moisture content; total fungi load initial =  $6.9 \times 10^4$ ; initial mc = 9.43%/ ND = not detected; <sup>c</sup> = concentration AFLs µg kg in duplicate; LOD 0.25; 0.08; 0.25 and 0.08 µg/kg e LOQ 0.50; 0.17; 0.50 and 0.17 µg/kg, to AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub> respectively; <sup>d</sup> AFB<sub>1</sub> + AFG<sub>1</sub> + AFB<sub>2</sub> + AFG<sub>2</sub>. Temperature: 20°C (16.6 – 20.6°C); r.h.: 80% (75.8 – 85)

#### 3.1.1 Total fungi load

As expected, the in-shell Brazil nuts ozonation showed to be effective on the fungi/spores destruction during the period of storage. A reduction of total fungi count was registered as soon as the first day after O<sub>3</sub> application in all treated nut Groups. However, the complete destruction of fungi (no growth: 'ng' label) was reached at different stages of storage depending upon the gas concentration applied. Total fungi destruction started at the first day when the highest O<sub>3</sub> concentration was applied (31.5 mg/L) and after 30 days of storage, when O<sub>3</sub> exposure was registered in all O<sub>3</sub> treated nut Groups after a month of storage up to the end of the six month. From the original (untreated nuts) total fungi load of 4.83 log cfu (Control), it went down at day one of O<sub>3</sub> treatment to 3.5 and 3.3 log cfu/g for groups I (O<sub>3</sub>: 10.0 mg/L) and II (O<sub>3</sub>: 14.0 mg/L), respectively; and no fungi growth was detected in the nuts treated with the highest O<sub>3</sub> concentration (Group III – O<sub>3</sub>: 31.5 mg/L). These figures were very different of those observed with the

nuts of control groups, where the fungi load remained somewhat stable with an insignificant increase during the whole period of storage i.e., from -zero/-one (4.83 log cfu/g) to 180 d (4.91 log cfu/g). Similar situation happened to those nuts when analyzed after being shelled, however in a lower extend from D-one (2.54 log cfu/g) to 180 d (2.69 log cfu/g) – control group) which was expected, as the edible part was protected by the shell. Some works have reported the effect of ozonation in food and nuts and they have similar findings for fungi load reduction as obtained in the present experiments, however applying lower O<sub>3</sub> concentrations. We utilized higher O<sub>3</sub> concentrations due to the fact that our aim was more than just fungi disinfection. We wanted to obtain a significant AFLs degradation too (to be discussed in the next Session) and for that purpose the O<sub>3</sub> concentration should be higher. Zhao and Cranston (1995) studied the effect of O<sub>3</sub> on black pepper at a lower gas concentration (6.7 mg/L) than in our study. The authors reported fungi load reduction from 7 to 4 log cfu/g. In another study carried out in dried figs (Oztekin et al., 2006), O<sub>3</sub> reduced the total fungi load from initial 1.46 to 1.00, 0.57, and 0.40 log cfu/g using O<sub>3</sub> at 1, 5, and 10 mg/L, respectively. In a work carried out on Brazil nut, however treating the nuts with aqueous O<sub>3</sub> solutions (0, 1, 10 and 20 mg/L) the authors showed that these treatments were effective to fungi control reaching an inactivation rate of 100% for *A. parasiticus* and 96% for *A. flavus*. However O<sub>3</sub> application in water can lead to an increase of moisture, reducing crunchy and firmness (Freitas and Venancio, 2008).

### 3.1.2. Aflatoxigenic species of *Aspergillus*

Regarding the isolation of aflatoxigenic species of *Aspergillus* (*A. flavus* and *A. parasiticus*) in the Brazil nuts just after the O<sub>3</sub> treatments (Group I, II, III), only the highest O<sub>3</sub> concentration treated nuts (Group III - 31.5 mg/L) did not allow them to grow since day one after gas application. The same occurred to the Groups of lower O<sub>3</sub> concentrations, but only after 30 d of storage. On the other hand, the Control Group was contaminated permanently by these species throughout the whole storage period. Similar situation happened when Mason et al. (1997) applied O<sub>3</sub> (5 mg/L) in cultures of *A. flavus* and *Fusarium verticillioides*. Ozone treatment in these conditions inhibited the growth, sporulation and mycotoxin production. Kells et al. (2001) studied the efficacy of O<sub>3</sub> as a fumigant to disinfest stored maize from insects. At a concentration of 50 mg/L, these authors observed a reduction of 63% of the kernel contamination level by *A. parasiticus*. It seems important to emphasize that the species *A. nomius* was not detected in the present experiments either due to the fact that after nut dehydration, fungi were destroyed by the heating temperature or fungi competition as nuts were purchased in the market in the South of Brazil, or also because the AFPA media does not give a clear response, i.e. not enhancing the characteristics of that *Aspergillus* species.

### 3.1.3. Environmental conditions after O<sub>3</sub> treatment: r.h. level

During the period of the in-shell Brazil nut storage in Southern Brazil were rather mild with average temperature of 20°C (min 16.6 and max 20.6°C) and r.h. of 80% (min 75.8 and max 85.3%).

### 3.1.4. Effect of O<sub>3</sub> on AFL contamination

In contrary to the nuts Control Group, a reduction on the AFL levels was detected throughout the whole storage period of the in-shell Brazil nuts O<sub>3</sub> treated (Table 1). Just after the O<sub>3</sub> treatments, either at gas concentration of 14 or 31.5 mg/L, the Brazil nuts did not present contamination by AFLs – up to the limit of quantification of the method used (0.50, 0.17, 0.50, and 0.17 µg/kg, for AFB<sub>1</sub>, AFB<sub>2</sub>, AFG<sub>1</sub> and AFG<sub>2</sub>, respectively). Although the three concentrations applied were able to degrade the toxins, some AFLs were still detected: 3.01 µg/kg (74% reduction) at the lower O<sub>3</sub> concentration (10 mg/L) after the first month of storage, the toxins was able to be totally degraded after 30 d. As far as the storage period and the AFL degradation are concerned, all the groups did not present any AFLs contamination after one month of storage onwards. As expected with the Control Group, the nuts AFL level remained stable or slightly increased from the beginning to the end of storage (11.58 to 12.55 µg/kg, respectively). That could be explained by the controlled experiment environment and storage conditions applied, and additionally, by the mild/low temperatures in Southern Brazil. The fact that other AFLs were detected, i.e., AFB<sub>2</sub> and AFG<sub>2</sub> in the Control Group, was probably the consequence of the origin of the nuts which were purchased in the retail market of Southern Brazil and not in the Amazon region, thus exposed to different fungi species contamination through manipulation and different environments, from tropical to temperate climate down in the south, respectively.

### 3.1.5. O<sub>3</sub> nuts sensory attributes for quality acceptance

No significant changes ( $P < 0.05$ ) were found between shell and nut appearances, strange odor, roasted flavor, rancidity and firmness scores of the ozonated Brazil nuts samples stored and they did not differ greatly among the concentrations used too. They were between 3 (indifferent) and 4 (like), different of the Control Group that received score 2 for most of the attributes except for nut firmness (score 3), showing that the treatment with O<sub>3</sub> and the period of storage of the in-shell Brazil nuts, did not affect their sensory quality attributes for all O<sub>3</sub> treated Groups. Also the shell received score 4, except for roasted flavor (score 3). Therefore, regarding sensory evaluation, the gas treatment kept nuts sensory attributes thus still palatable apart from being safer.

Prevention is a better strategy than detoxification which is much more complicated and so the implications to human and animal health. Despite of the findings, there is a need of more studies, especially on application in pilot plants utilizing larger amounts of nuts under the Amazon forest environment (first and second storages) prior factory processing, in order to establish the optimal applicable O<sub>3</sub> gas concentration and time of exposure for maximum effectiveness utilizing the present findings. This work is being applied in a pilot plant at a Brazil nut factory in the Amazon State for checking its effectiveness under the Amazon region environment. Important to emphasize that gaseous O<sub>3</sub> decomposes to form O<sub>2</sub> and it does not affect the environment, nor leave residues in the nuts.

#### Acknowledgements

This work is part of a Research Project on “Technology Development for Prevention and Control of AFLs in Stored Brazil Nuts” that has been developed in the Food and Technology Department of UFSC/SC, Brazil.

#### References

- AOAC, 2005. Official Methods of Analysis International Nuts and Nuts Products., Art. 925.40, 18<sup>th</sup> ed. Vol II, chapter 40.
- APHA - American Public Health Association, 1980. Standard methods for the examination of water and wastewater. 15 ed. Washington: American Public Health Association.
- Arrus, K., Blank, G., Abramson, D., Clear, R., Holley, R.A., 2005. AFL production by *Aspergillus flavus* in Brazil nuts. *Journal of Stored Products Research* 41, 513–527.
- CONFORCAST, 2008. Plano de amostragem para castanha-do-Brasil. Access: 02/06/2008, //www.nutfruit.org.
- De-Mello, F.R.; Scussel, V.M., 2007. Characteristics of in-shell Brazil nuts and their relationship to aflatoxin contamination: criteria for sorting. *Journal of Agriculture and Food Chemistry* 55, 9305-9310.
- De-Mello, F.R.; Scussel, V.M. 2009. Development of physical and optical methods for In-shell Brasil nuts sorting and aflatoxin reduction. *Journal of Agricultural Science*, 1, 3-14.
- ECC – European Communities Commission, 2006. – Commission Regulation 1881/2006 of 19 Dec 2006 setting maximum levels for certain contaminants in foodstuff. *Official Journal of European Communities*, L.364/5-364/24.
- Frazier, W. C., Westhoff, D.C., 1988. *Food Microbiology*. McGraw-Hill Book Company, 4<sup>th</sup> ed, New York.
- Freitas, O., Venancio, A., 2008. Gaseous ozone application on Brasil nuts. Encontro Brasileiro de Micotoxinas, julho, Rio de Janeiro, RJ. poster 24.
- Inan F., Pala, M., Doymaz, I., 2007. Use of ozone in detoxification of aflatoxin B1 in red pepper. *Journal of Stored Products Research* 43, 425-429.
- Kells, A.A., Mason, L.J., Maier, D.E., Woloshuk, C.P., 2001. Efficacy and fumigation characteristics of ozone in stored maize. *Journal of Stored Products Research* 37, 371–382.
- Machida, S., Saito, M., 1999. A rapid identification method for aflatoxin-producing strains of *Aspergillus flavus* and *A. parasiticus* by ammonia vapor. *Mycoscience* 40, 205-208.
- Maeba, H., Takamoto, Y., Kamimura, M., Miura, T., 1988. Destruction and detoxification of aflatoxins with ozone. *Journal of Food Science* 53, 667–668.
- Mangan, N. and Aldred, 2007. Post-harvest control strategies: minimizing mycotoxins in the food chain. *International Journal of Food Microbiology* 119, 131-139.
- Mason, L.J., Woloshuk, C.P., Maier, D.E., 1997. Efficacy of ozone to control insects, molds and mycotoxins. In: Donahaye, E.J., Navarro, S. & Varnava, A., (Eds), *Proceedings of the International Conference on Controlled Atmosphere & Fumigation in Stored Products*, Cyprus Printer Ltd., Nicosia, pp. 665–670.

- McKenzie, K.S., Sarr, A.B., Maymura, K., Bailey, R.H., Miller, D.R., Rogers, T.D., Norred, W.P., Voss, K.A., Plattner, R.D., Kubena, L.F., Phillips, T.D., 1997. Oxidative degradation and detoxification of mycotoxins using a novel source of ozone. *Food and Chemical Toxicology* 35, 807-820.
- Olmez, H., Akbas, M.Y., 2009. Optimization of ozone treatment of fresh-cut green leaf lettuce. *Journal of Food Engineering* 90, 489-494.
- Olsen, M., Johsson, P., Moller, T., Paladino, R., Lindblat, M., 2008. Is *Aspergillus nomius*, an important aflatoxin producer in Brazil nuts? *World Mycotoxin Journal* 1, 123-126.
- Oztekin, S., Zorlugenç, B., Zorlugenç, F.K., 2006. Effects of ozone treatment on microflora of dried figs. *Journal of Food Engineering* 75, 396-399.
- Pacheco, A.M., Scussel, V.M., 2009. Aflatoxins evaluation on in-shell and shelled dry Brazil nuts for export analysed by LC-MS/MS-2006 and 2007 harvests. *World Mycotoxin Journal* 2, 295-304.
- Pacheco, A.M., Scussel, V.M., 2007. Selenium and aflatoxin levels in raw Brazil nuts from the Amazon basin. *Journal of Agriculture and Food Chemistry* 55, 11087-11092.
- Pacheco, A.M., Scussel, V.M., 2006. Castanha do Brasil – da Floresta ao Consumidor. Ed. Editograf, Brazil.
- Perez, A. G., Sanz, C., Rios, J. J., Olias, R., Olias, J. M., 1999. Effects of ozone treatment on postharvest strawberry quality. *Journal of Science of Food and Agriculture* 47, 1652-1656.
- Pitt, J.I., Hocking, A.D., 1997. *Fungi and food spoilage*. 2<sup>nd</sup> ed. London: Blackie Academic & Professional, London.
- Pitt, J.I., Hocking, A.D., Glenn, D.R., 1983. An improved medium for the detection of *Aspergillus flavus* and *A.parasiticus*. *Journal of Applied Bacteriology* 54, 109-114.
- Shamarajeewa, U., Sen, A.C., Cohen, M.D., Wei, C.I., 1990. Detoxification of aflatoxins in foods and feeds by physical, chemical methods. *Journal of Food Protection* 53, 489-501.
- Samsom, R.A., Hoeskra, E.S., Frisvad, J.C., 2004. *Introduction to Food and Airborne Fungi*. 7<sup>th</sup> edition, CBS. The Netherlands.
- Sobolev, V.S., 2007. Simple, rapid, and inexpensive cleanup method for quantitation of aflatoxins in important agricultural products by HPLC. *Journal of Agriculture and Food Chemistry* 55, 2136-2141.
- Stone, H.S., Sidel, J.L., 1993. *Sensory Evaluation Practices*. Academic Press, New York.
- Yesilcimen, A.M., Murat, O., 2006. Effect of treatments on AFLs degradation & physicochemical properties of pistachios. *Journal of Science of Food and Agriculture* 86, 13, 2099-2104.
- Zhao, J., Cranston, P.M., 1995. Microbial descontamination of black pepper by ozone & effects of treatment on volatile oil constitutes of spices. *Journal of Science of Food and Agriculture* 68, 11-18.