

- Belton, J.M., McCord, R.P., Gibcus, J.H., Naumova, N., Zhan, Y. and Dekker, J., 2012: Hi-C: A comprehensive technique to capture the conformation of genomes. *Methods* 58, 268-276.
- Burton, J.N., Adey, A., Patwardhan, R.P., Qiu, R., Kitzman, J.O. and Shendure, J., 2013: Chromosome-scale scaffolding of de novo genome assemblies based on chromatin interactions. *Nature Biotechnology* 31, 1119.
- Chesters, D., Zheng, W.M. and Zhu, C.D., 2015: A DNA barcoding system integrating multigene sequence data. *Method in Ecology and Evolution* 6, 930-937.
- Dönitz, J., Schmitt-Engel, C., Grossmann, D., Gerischer, L., Tech, M., Schoppmeier, M., Klingler, M. and Bucher, G., 2014: iBeetle-base: A database for RNAi phenotypes in the red flour beetle *Tribolium castaneum*. *Nucleic Acids Research* 43, D720-725.
- Futahashi, R., Tanaka, K., Matsuura, Y., Tanahashi, M., Kikuchi, Y. and Fukatsu, T., 2011: Laccase2 is required for cuticle pigmentation in stinkbugs. *Insect Biochemistry and Molecular Biology* 41, 191-196.
- Gurevich, A., Saveliev, V., Vyahhi, N. and Tesler, G., 2013: QUAST: Quality assessment tool for genome assemblies. *Bioinformatics* 29, 1072-1075.
- Li, H. and Durbin, R., 2009: Fast and accurate short read alignment with burrows-wheeler transform. *Bioinformatics* 25, 1754-1760.
- McKenna, D.D., Scully, E.D., Pauchet, Y., Hoover, K., Kirsch, R., Geib, S.M., et al., 2016: Genome of the Asian longhorned beetle (*Anoplophora glabripennis*) reveals key functional and evolutionary innovations at the beetle-plant interface. *Genome Biology* 17, 227.
- Rastas, P., Calboli, F.C., Guo, B., Shikano, T., and Merila, J., 2015: Construction of ultradense linkage maps with Lep-MAP2: stickleback F2 recombinant crosses as an example. *Genome Biology and Evolution* 8, 78-93.
- Schlipalius, D.I., Valmas, N., Tuck, A.G., Jagadeesan, R., Ma, L., Kaur, R., Goldinger, A., Anderson, C., Kuang, J., Zury, S. and Mau, Y.S., 2012: A core metabolic enzyme mediates resistance to phosphine gas. *Science* 338, 807-810.
- Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V. and Zdobnov, E.M., 2015: BUSCO: On the proper publication of stored product protection research – conference proceedings are better than you thought. BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31, 3210-3212.
- Talla, V., Suh, A., Kalsoom, F., Dincă, V., Vila, R., Friberg, M., Wiklund, C. and Backström, N., 2017: Rapid increase in genome size as a consequence of transposable element hyperactivity in wood-white (*Leptidea*) butterflies. *Genome Biology and Evolution* 9, 2491-2505.
- Wieslander, L., 1979: A simple method to recover intact high molecular weight RNA and DNA after electrophoretic separate in low gelling temperature agarose gels. *Analytical Biochemistry* The control, 305-309.
- Zheng, X.H., Lu, F., Wang, Z.Y., Zhong, F., Hoover, J. and Mural, R., 2004: Using shared genomic synteny and shared protein functions to enhance the identification of orthologous gene pairs. *Bioinformatics* 21, 703-710.

DNA barcode of stored-product Pests based on Mitochondrial Cytochrome Oxidase I Gene

Yi Wu^{1*}, Zhihong Li², Fujun Li¹, Václav Stejskal³, Dan Zheng¹, Xin Chen¹, Yang Cao¹

¹Academy of State Administration of Grain, No. 11 Baiwanzhuang Street, Beijing 100037, China.

²Department of Entomology, College of Agronomy and Biotechnology, China Agricultural University, Yuanmingyuan West Rd. 2, Haidian District, Beijing 100193, China.

³Department of Pest Control of Stored Products and Food Safety, Crop Research Institute, Drnovská 507, Prague, Czech Republic.

*Corresponding author: wuyi@chinagrains.org

DOI 10.5073/jka.2018.463.028

Abstract

The stored-product pests are economically important that can be spread through grain trade. Most stored-product pests, including eggs, nymphs, and adults, are very small and difficult to identify morphologically. Also the classification and identification of them have always been hindered by the overwhelming number of species, widely distribution. Here, we collected 43 stored-product pests from 46 geographical locations in China and other countries. The mtDNA COI gene sequences were sequenced. Software MEGA 5 was used to analyze the sequence comploition and genetic distances. Three molecular phylogenetic trees of Platypodidae were recomstruced using PAUP4.0 according to distance/ the neighbour-joining (NJ) and maximum parsimony (MP). The molecular results were compared with the morphological taxonomy. The interspecific genetic distance of the stored-product pests was significantly higher than the intraspecific genetic distance according to the barcoding gap analysis. This work provides a practical approach for the precise and rapid diagnosis of stored-product pests.

Keywords: DNA Barcoding, stored-product pests, mtDNA COI gene, phylogeny

Introduction

Stored-grain insects and mites are great economic significance for grains and other products during storage. They are closely related with human life. The rapid identification is the precondition and basis for the comprehensive prevention and control of the pests. The modern identification of stored-grain insects by molecular biology techniques is able to get rid of the influence of the growth situation of individual specimen and the environment, and get accurate and reliable result from their DNA.

Traditionally, the species have been identified based on morphological characteristics of the adult. However, the identification of the species based on immature stages (i.e., egg, larva or pupa) or adult body parts, which lack distinct diagnostic characteristics, is very difficult and sometimes not reliable (Li et al., 2011). Traditional morphological identification is also time-consuming, requires specialized taxonomic knowledge and microscopy techniques (Yang et al., 2013; Jiang et al., 2014)

Recently, molecular identification based on nucleotide sequence analysis has become an effective method used to complement traditional taxonomic identification. For some important insect pests of stored products, AFLP has been used to diagnostic *Liposcelis* (Qin et al., 2008; Li et al., 2011), *Sitophilus oryzae* and *Sitophilus zeamais* (Hidayat et al., 1996), DNA barcode technology for *Liposcelis entomophila* (Yang et al., 2012). Some recent studies have implemented by PCR with species-specific primer pairs (Zhao et al., 2016). Species-specific primer identification is a PCR-based procedure that yields a unique band of known size and allows a species to be identified directly after gel electrophoresis (Wu et al., 2016). In animals, mitochondrial cytochrome c oxidase I (COI) gene, has been shown to be a reliable, quick and cost-effective tool for the identification of organisms of various taxa in all life stages. More and more important insect pests have been identified by this way (Namikoshi et al., 2011; Zhang et al., 2012; Jiang et al., 2014). It is a great advantage especially for the identification of small size pests.

DNA barcoding is a DNA-based species identification system which offers a promising supplemental technique with standardized portions of the genome (Hebert et al., 2003). The most commonly used barcode gene, mitochondrial (mt) DNA cytochrome c oxidase I (COI), has been shown to be a reliable, quick and cost-effective tool for the identification of organisms of various taxa in all life stages (Augot et al., 2010; Cywinska et al., 2010). A threshold of 2-3% mtDNA COI sequence divergence was recommended to define separate species for insects and mammals (Hebert et al., 2003). In studies of butterflies and ants, DNA barcoding has been successful in defining species boundaries by genetic distance thresholds (Hebert et al., 2004; Smith et al., 2005); however, there is no established universal distance threshold value to distinguish between taxonomic groups.

In the present study, we describe a reliable and efficient method based on conventional PCR with mtDNA COI gene, and we set up a DNA barcode data base for stored-product Pests, which we hope will prove useful for the rapid diagnosis.

Materials and Methods

Specimens used in this study are collected from different provinces in China, including 46 geographical locations as fig. 1, some mites from Czech Republic, and except *Liposcelis bostrychophila*, *L. entomophila*, *L. decolor*, and *L. paeta*, the samples of *Liposcelis* from the Plant quarantine laboratory of China Agricultural University (CAUPQL). There are 43 species of stored-grain insects/ mites in total, 415 individuals, every species at least 5 specimens.

Total genomic DNA was extracted from the entire body of individual adults using the TIANamp Genomic DNA kit (DP304, TIANGEN, China) following the manufacturer's protocol for animal tissue. Five individuals from each species were used. PCR was performed with a pair of universal primers, LCO1490 (fw) 50 GGT CAA CAA ATC ATA AAG ATA TTG G 30 and HCO2198 (rev) 50 TAA ACT TCA GGG TGA CCA AAA AAT CA 30, amplifying an approximately 710 bp fragment of the standard mtDNA COI-5 barcode (Folmer et al., 1994). PCR products were separated on a 1.0% (w/v) agarose gel (1 × TAE), stained with ethidium bromide, and visualised under UV light. The agarose gel slice

containing the PCR amplicon of interest was excised and placed in a centrifuge tube. The agarose gel slice containing the PCR amplicon of interest was excised and the DNA was gel extracted. Bidirectional sequencing reactions were carried out from a single individual of each geographical isolate (Beijing Aoke Biotechnology Co., Ltd.).

DNAMAN software (Lynnon Biosoft, Vaudreuil, Quebec, Canada) was used for DNA multiple sequence alignment using an optimal alignment method. Genetic diversity was estimated for haplotype diversity (Hd) and nucleotide diversity in DnaSP version 4.10.1 (Librado & Rozas, 2009). Pairwise genetic distances for COI were calculated using the Kimura-2-Parameter (K2P) distance model implemented in the software Molecular Evolutionary Genetics Analysis 5 (MEGA 5; Tamura et al., 2011). All phylogenetic analyses were carried out using the program PAUP 4.0 (Swofford, 2002). Two different types of phylogenetic trees, neighbour-joining (NJ) and maximum parsimony (MP), were graphically displayed and compared. A heuristic search was employed using tree bisection and reconnection (TBR) branch swapping and random addition for 100 replicates, and bootstrapping was performed using 1000 replications.

Results

In this study, we tested and evaluated the general genes of COI, which is an appropriate gene for identifying the DNA barcode of stored-grain insects/mites – a section of 650bp COI gene in mitochondria (primer pair LCO490/HCO2198). The mtDNA COI sequences of 415 obtained in this study. The sequences were all trimmed to a 650 bp core region that could be unambiguously aligned to one another. No sequences contained indels or nonsense codons, allowing for easy alignment and supporting their origin in the mitochondrial gene.



Fig. 1 Distribution of sampling sites for stored grain insects and mites in China

A rapid identification DNA barcode sequence database of stored-grain insect/mite is established, including 43 species of stored-grain insects/mites. Every species has at least 5 specimens, we get the COI barcode sequence of 415 specimens. 98 haplotypes, among which only 1 haplotype is found in 20 species of insects/mites, showing the diversity of stored-grain insect is relatively low. Also most showed low intra-species divergence. Based on DNA barcode sequence and distance method, the neighbour-joining (NJ) and maximum parsimony (MP), the phylogenetic tree of stored-grain insect/mite is built. Both the NJ and MP phylogenetic analysis of the COI gene generated the same tree topology. The resulting trees showed a clear clade and every species has individual branch.

DNA barcode in the mitochondria COI sequence can be used to identify the species of stored-grain insects/mites rapidly and accurately.




Fig. 2 Neighbour-joining tree of stored product insects and mites

Acknowledgement

Financial support for this research was provided by National Key R&D Program of China (2016YFD0401004-2), the Special Fund for Grain Scientific Research in the Public Interest of China (201513002).

References

- Augot, D., Sauvage, F., Joue, D., Simphal, E., Veuille, M., Couloux, A., Kaltenbach, M.L., Depaquit, J., 2010: Discrimination of *Culicoides obsoletus* and *Culicoides scoticus*, potential bluetongue vectors, by morphometrical and mitochondrial cytochrome oxidase subunit I analysis. *Infection, Genetics and Evolution* 10, 629-637.
- Cywinska, A., Hannan, M.A., Kevan, P.G., Roughley, R.E., Iranpour, M., Hunter, F.F., 2010: Evaluation of DNA barcoding and identification of new haplomorphs in Canadian deerflies and horseflies. *Medical and Veterinary Entomology* 24, 382-410.
- Folmer, O., Black, M., Hoeh, W., Lutz, R., Vrijenhoek, R., 1994: DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3, 294-299.
- Hebert, P.D.N., Cywinska, A., Ball, S.L., DeWaard, J.R., 2003: Biological identifications through DNA barcodes. *Proceedings of the Royal Society B* 270, 313-321.
- Hebert, P.D.N., Stoeckle, M.Y., Zemlak, T.S., Francis, C.M., 2004: Identification of birds through DNA barcodes. *PLoS Biology* 2, 1657-1663.
- Hidayat, P., Phillips, T.W., Ffrench-Constant, R.H., 1996: Molecular and morphological characters discriminate *Sitophilus oryzae* and *S. zeamais* (Coleoptera: Curculionidae) and confirm reproductive isolation. *Annals of the Entomological Society of America* 89, 645-652.
- Jiang, F., Li, Z.H., Wu, J.J., Wang, F.X., Xiong, H.L., 2014: A rapid diagnostic tool for two species of *Tetradacus* (Diptera: Tephritidae: Bactrocera) based on species-specific PCR. *Journal of Applied Entomology* 6, 418-422.
- Li, Z.H., Kučerová, Z., Zhao, S., Stejskal, V., Opit, G., Qin, M., 2011: Morphological and molecular identification of three geographical populations of the storage pest *Liposcelis bostrychophila* (Psocoptera). *Journal of Stored Products Research* 47, 168-172.
- Librado, P. and Rozas, J., 2009: DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, 25, 1451-1452.
- Namikoshi, A., Takashima, Y., Iguchi, J., Yanagimoto, T., Yamashita, M., 2011: Species identification of Alaska pollock, *Gadus* spp., and *Micromesistius* spp. in cod roe products using a PCR-based method. *Fisheries Science* 77, 671-678.
- Qin, M., Li, Z., Kučerová, Z., Cao, Y., Stejskal, V., 2008: Rapid discrimination of the common species of the stored product pest *Liposcelis* (Psocoptera: Liposcelididae) from China and the Czech Republic based on PCR-RFLP analysis. *European Journal of Entomology* 105, 713-717.
- Smith, M.A., Fisher, B.L., Hebert, P.D.N., 2005: DNA barcoding for effective biodiversity assessment of a hyperdiverse arthropod group: the ants of Madagascar. *Proceedings of the Royal Society B* 360, 1825-1834.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011: MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*. doi:10.1093/molbev/msr121.
- Wu, Y., Li, F., Li, Z., Stejskal, V., Aulicky, R., Kučerová, Z., Zhang, T., He, P., Cao, Y., 2016: Rapid diagnosis of two common stored-product predatory mite species based on species-specific PCR. *Journal of Stored Products Research* 69, 213-216.
- Yang, Q., Kučerová, Z., Li, Z., Kalinović, I., Stejskal, V., Opit, G., Cao, Y., 2012: Diagnosis of *Liposcelis entomophila* (Insecta: Psocodea: Liposcelididae) based on morphological characteristics and DNA barcodes. *Journal of Stored Products Research* 48, 120-125.
- Zhang, G.F., Meng, X.Q., Min, L., Qiao, W.N., Wan, F.H., 2012: Rapid diagnosis of the invasive species, *frankliniella occidentalis*, (Pergande): a species-specific COI marker. *Journal of Applied Entomology* 136, 410-420.
- Zhao, Z.H., Cui, B.Y., Li, Z.H., Jiang, F., Yang, Q.Q., Kučerová, Z., Stejskal, V., Opit, G., Cao, Y., Li, F.J., 2016: The establishment of species-specific primers for the molecular identification of ten stored-product psocids based on ITS2 rDNA. *Scientific Reports* 6, 210-222.

Effect of delayed mating on reproductive performance of *Lasioderma serricorne* (F.) (Coleoptera: Anobiidae)

Rizana Mahroof^{1*}, Barbara Amoah¹, Alison Gerken², Jim Campbell²

¹Department of Biological and Physical Sciences, South Carolina State University, Orangeburg, South Carolina, 29117, USA

²USDA, Agricultural Research Service, Center for Grain and Animal Health Research, Manhattan, Kansas, 66502, USA

*Corresponding Author: rmahroof@scsu.edu

DOI 10.5073/jka.2018.463.029