

- [15] Kacimi El Hassani A, Dacher M, Gary V, Lambin M, Gauthier, M and Armangaud, C: Effects of sublethal doses of acetamiprid and thiamethoxam on the behaviour of the honeybee (*Apis mellifera*), Arch Environ Contamin Toxicol, **54**(4): 653-61(2008)
- [16] Guez D, Suchail S, Gauthier M, Maleszka R and Belzunces LP: Contrasting effects of imidacloprid on habituation in 7- and 8- days-old honeybees, Neurobiology of Learning and Memory, **76** (2), 183 – 191 (2001)
- [17] Decourtye A, Lacassie E and Pham-Delègue MH: Learning performances of honeybees (*Apis mellifera* L) are differentially affected by imidacloprid according to the season, Pest Manag Sci **59**: 269–278 (2003)
- [18] Aupinel P, Fortini D, Michaud B, Marolleau F, Tasei JN and Odoux JF: Toxicity of dimethoate and fenoxycarb to honey bee brood (*Apis mellifera*) using a new in vitro standardized feeding method, Pest Manag Sci **63**: 1090 – 1094
- [19] Vandame R, Meled M, Colin ME and Belzunces LP: Alteration of the homing-flight in the honey bee *Apis mellifera* L. exposed to sublethal dose of deltamethrin. Environ. Toxicol. Chem. **14**:855-860 (1995)
- [20] Decourtye A, Devillers J, Cluzeau S, Charreton M and Pham-Delègue MH: Effect of imidacloprid and deltamethrin on associative learning in honeybees under semi-field and laboratory conditions, Ecotoxicology and Environmental Safety, **57**, 410 – 419 (2004)
- [21] Cuthbertson AG, Walters KF and Deppe C: Compatibility of the entomopathogenic fungus *Lecanicillium muscarium* and insecticides for eradication of sweetpotato whitefly, *Bemisia tabaci*. Mycopathologia. **Aug.160v(1)**: 35-41 (2005)
- [22] Feng MG and Pu XY: Time–concentration–mortality modeling of the synergistic interaction of *Beauveria bassiana* and imidacloprid against *Nilaparvata lugens*, Pest Manag Sci **61**:363–370 (2005)
- [23] Santos AV, Lorenz de Oliveira B and Samuels RI: Selection of entomopathogenic fungi for use in combination with sub-lethal doses of imidacloprid: perspectives for the control of the leaf-cutting ant *Atta sexdens rubropilosa* Forel (Hymenoptera: Formicidae), Mycopathologia **163**:233–240 (2007)
- [24] Vandame R and Belzunces LP: Joint actions of deltamethrin and azole fungicides on honey bee thermoregulation, Neuroscience Letters **251** (issue1): 57-60 (1998)

III. Bumblebees and other bee species

The impact of different concentrations of a pyrethroid insecticide on the cyclic gas exchange cycles on bumble bees

*Karise, R., Kuusik, A., Mänd, M., Metspalu, L., Hiisaar, K., Luik, A.

Institute of Agricultural and Environmental Sciences, Estonian University of Life Sciences, 1 Kreutzwaldi St., 51014 Tartu, Estonia. *Reet Karise, PhD, reet.karise@emu.ee

Abstract

Minor effects of pesticides may remain unnoticeable in adult bees because of no visible changes in their behaviour throughout several days after coming into contact with pesticides. The hypothesis of this work is that changes which are not observable through the behaviour of the bumble bees can be seen through physiological patterns. The aim of the present research was to study the effect of low concentrations of Fastac 100 EC on discontinuous gas exchange cycles of bumble bee *Bombus terrestris* foragers. Using a system of flow-through CO₂ respirometry, the effect of different concentrations of alpha-cypermethrin on bumble bee foragers was studied. We found that the concentration of Fastac 100 EC that is used in the fields and a tenfold solution of that caused significant decrease in the frequency of bursts of CO₂ releases in bumble bees. 20-fold diluted solution did not cause the significant decrease. The lifespan of treated bumble bees also decreased by the field concentration and ten-fold diluted concentration. Alpha-cypermethrin caused changes in the respiration patterns of *B. terrestris* foragers although not always seen through the behaviour. These changes could potentially lead to a decreasing individual and colony survival.

Keywords: Respiration cycles, pyrethroid insecticide, bumble bees