

## Karyotype studies on grape phylloxera (*Daktulosphaira vitifoliae* FITCH)

by

ASTRID FORNECK<sup>1)</sup>, Y. JIN<sup>2)</sup>, A. WALKER<sup>2)</sup> and R. BLAICH<sup>1)</sup>

<sup>1)</sup> Universität Hohenheim, Institut für Obst-, Gemüse- und Weinbau, Fachgebiet Weinbau (370), Stuttgart, Deutschland

<sup>2)</sup> University of California, Davis, Department of Viticulture and Enology, Davis, USA

**S u m m a r y :** A cytogenetic technique was developed to produce suitable chromosome spreads for phylloxera karyotype analysis. The karyotype for pathogenetic phylloxera was  $2n = 10$ . Karyotypes from haploid sex cells were found to vary between  $n = 5$  and  $n = 6$ , the latter possibly indicating an aneuploidic aberration. Tetra- and polyploid cells were detected in somatic trophocytes. Preparation of phylloxera sex and somatic cells for chromosomal analysis reported here enables the study of genetic variation on a chromosomal scale.

**Key words :** chromosome, *Daktulosphaira vitifoliae*, karyotype, phylloxera.

### Introduction

Phylloxera, *Daktulosphaira vitifoliae* FITCH, is one of the most important grape pests. Their importation and subsequent spread on the susceptible roots of European vineyards in the last half of the 19th century led to the development of hundreds of resistant rootstocks upon which most of the world's vineyards are now planted. Phylloxera reproduce through cyclical parthenogenesis on grape roots and leaves. Populations with a complete life cycle (holocyclic) bridge these parthenogenetic stages with dimorphic forms that produce sexual offspring (*sexuparae*). These offspring mate and lay an overwintering egg that initiates another parthenogenetic cycle (the *fundatrix*). Populations that only reproduce parthenogenetically are termed anholocyclic.

Phylloxera resistant rootstocks have successfully controlled this pest for over 100 years. However, strains with more aggressive feeding habits have been reported in Germany (BOUBALS 1994) and California (GRANETT *et al.* 1987). Studies on the population dynamics and biology of these and other phylloxera revealed the occurrence of new phylloxera biotypes that are host plant-adapted. These discoveries led to descriptions of genetic variation within and among phylloxera populations in North America and Europe (FONG *et al.* 1995; FORNECK *et al.* 1998; LIN *et al.* 1999). Efforts to associate phylloxera feeding behavior with a distinct genotype, or genetic marker have not yet been successful. Genetic markers linked to aggressive feeding behavior would be useful in pest management and rootstock breeding. It might also be possible to associate karyotypes or levels of aneuploidy with changes in phylloxera behavior. Such chromosomal changes have been found in aphids and have been associated with the evolution of insecticide resistance (BLACKMAN and TAKADA 1975; BLACKMAN 1987). The objective of the study presented here

was to define the karyotypes of 6 phylloxera populations from Germany, France, Italy and California (USA). This process required the development and modification of cytogenetic techniques, and will aid future investigations into the genetic basis of phylloxera behavior and biotype development.

### Material and Methods

**Phylloxera material:** Phylloxera originating from different European sources (Table) were raised on grape host plants using isolation chambers under controlled greenhouse conditions (16 h photoperiod, 24–26 °C temperature). Under these conditions parthenogenetic and sexual phylloxera morphs were produced.

**Chromosome preparations:** Ovarioles, pre-oviposited eggs and sexual larvae were dissected from anholocyclic adults or *sexuparae* in Ringer's saline solution for later examination of mitotic and meiotic cell divisions. These tissues were kept in a 1 % hypotonic solution of sodium citrate for about 15 min. Ten to 15 reproductive systems were pooled, transferred into a 500 µl Eppendorf tube, and fixed with three parts of acetic glacial acid, one part of methanol (abs.) for 10 min. After centrifuging (5 min at 500–800 g) the supernatant was poured off and the pellet resuspended in fresh fixative. The ovary tissue was macerated passing the suspension through a syringe 3 to 5 times. Cells were spread by dropping one or two drops of the suspension on a clean microscope slide, followed by air-drying. Chromosomes were stained with DAPI (4',6-diamino-2-phenylindole: 2 µg·ml<sup>-1</sup> in McIlvaine's buffer, pH 7) after a protocol published by MANICARDI *et al.* (1997) with minor modification, air-dried and examined through epifluorescent filter sets (Zeiss 01). For whole body preparations

Table

Populations and specimens used to examine phylloxera chromosomes. All populations were maintained under greenhouse culture on their original hosts

| Geographical origin       | Original host plant | Specimen analyzed   |
|---------------------------|---------------------|---|
| Bingen/Germany            | Teleki 5C           | Parthenogenetic ovaries   |
| Gundelsheim/Germany       | Teleki 5C           | Parthenogenetic ovaries   |
| Caprino di Veronese/Italy | Teleki 5C           | Parthenogenetic ovaries and eggs, ovaries from <i>sexuparae</i> , sexual larvae |
| Montpellier/France        | C3309               | Parthenogenetic ovaries and eggs  |
| California/USA            | Cabernet Sauvignon  | Parthenogenetic ovaries and eggs, ovaries from <i>sexuparae</i> , sexual larvae |

adult parthenogenetic phylloxera were placed on a microscope slide, mounted in a drop of dd H<sub>2</sub>O, squashed with a coverslip, and stained with a 0.01 % 3,6-bis (dimethyl-amino) acridine solution and examined as described above.

### Results and Discussion

**Phylloxera female reproductive system:** Phylloxera ovarioles are telotrophic like those commonly found in Hemiptera (Fig. 1 A, B). The number of ovarioles per adult female ranges from 9 to over 50 in fundatrices (females that initiate the first parthenogenetic generation of the season), the number being influenced by host and seasonal time (MAILLET 1957 b).




Fig. 1: Dissection of reproductive system of an adult parthenogenetic phylloxera in dorsal view. **A** Vital squash preparation prior to staining; **B** Stained with acridine orange. g: germarium; p: phylloxera; tf: terminal filament; ch: chorion. Bar represents 100  $\mu$ m.

**Karyotypes:** Karyotypes of parthenogenetic females from all 1998 collections were  $2n = 10$  (Fig. 2). Phylloxera chromosomes are similar to those of other Hemiptera; they are holocentric with no localized centromer. This makes condensed chromosomes in late prophase and metaphase appear as simple rods. Four pairs of autosomes and one pair of sex chromosomes could be observed, as reported in previous studies (MAILLET 1957 a). These 10 chromosomes could be divided into two classes: one pair of large chromosomes comprising an appendix similar to a chromosomal satellite, and 4 pairs of smaller chromosomes.

Endomeiotic recombination (COGNETTI 1961) was not observed during the mitotic division cycle in parthenoge-




Fig. 2: Karyotype for parthenogenetic phylloxera ( $2n = 8 + XX$ ).

netic phylloxera (Fig. 3 A, B). One karyotype containing  $2n = 9$  chromosomes was found in somatic cells of what appeared to be male sexual phylloxera. MAILLET (1957 a) described phylloxera sex cells and found ova with  $n = 5$  chromosomes and spermatids with  $n = 4$ . Spermatids were not observed but ova with  $n = 5$  and  $n = 6$  chromosomes were frequently found (Fig. 3 C, D). These cells had one additional chromosome that paired with one of the smaller autosomes. Supernumerary chromosomes have been described in aphids where they are termed B-chromosomes and do not affect the phenotype (BLACKMAN 1976). These B-chromosomes are usually shorter than A-chromosomes and do not pair with the standard chromosomes. The extra chromosome that we detected in ova cells did not meet the requirements in regard to morphology for B-chromosomes and thus the role of these aneuploidic cells remains unclear and requires further work.

Tetra- and polyploid cells derived from endomitosis were found in ovary cells of parthenogenetic females and are likely trophocytes that nurture developing oocytes (Fig. 3 E). Endopolyploidy is a common occurrence in aphids and widespread in somatic tissues like salivary glands, fat body tissue or trophocytes of oviparous females (BLACKMAN 1987). Multiple crossing over events were observed during diakinesis of meiosis II (Fig. 3 F).

### Conclusion

The cytogenetic techniques utilized here produced suitable chromosome spreads for karyotype analysis. The karyotype for parthenogenetic phylloxera was  $2n = 10$ . Karyotypes from haploid sex cells were found to vary between  $n = 5$  and  $n = 6$ . Endomeiosis was not observed. Polyploid cells were




Fig. 3: Phylloxera chromosomes displaying various karyotypes (A-F). **A** Partial mitotic cell division cycle of phylloxera: prometaphase, metaphase, early and late telophase; **B** Karyotype of a parthenogenetic female ( $2n = 10$ ); **C** Karyotype of a sexual phylloxera's ovum ( $n = 4 + X$ ); **D** Karyotype of a sexual phylloxera's ovum with additional chromosome ( $n = 4 + A + X$ ); **E** Tetraploid cell derived from trophocytes; **F** Multiple chiasmata in diakinesis during meiosis in a sexual phylloxera ovum. Bar represents 100  $\mu\text{m}$ .

detected in somatic trophocytes. The successful preparation of phylloxera sex and somatic cells for chromosomal analysis reported here is now being used to study genetic variation and its association with feeding behavior on a chromosomal scale.

### Acknowledgements

Financial support of the California Grape Rootstock Improvement Commission, the California Table Grape Commission, the American Vineyard Foundation, the German Research Society (Deutsche Forschungsgemeinschaft) and the German Academic Exchange Service (DAAD) is gratefully acknowledged.

### References

- BLACKMAN, R. L.; 1976: Cytogenetics of two species of *Euroraphis* (Homoptera, Aphididae). *Chromosoma* **56**, 393-408.
- -; 1987: Reproduction, cytogenetics and development. In: A. K. MINKS, P. HARREWIJN (Eds.): *Aphids, their Biology, Natural Enemies and Control*. 2B. Elsevier Amsterdam.
- -; TAKADA, H.; 1975: A naturally occurring chromosomal translocation in *Myzus persicae* (SULZER). *J. Entomol. (A)* **50**, 147-156.
- BOUBALS, D.; 1994: Phylloxéra, attention! En Allemagne, une race de phylloxéra radicicole plus agressive attaquerait les racines de SO 4, 5 BB, 5 C et 125 AA (tous *riparia x berlandieri*)! *Progr. Agric. Vitic.* **111** (17), 374-375.
- COGNETTI, G.; 1961: Endomeiosis in parthenogenetic lines of aphids. *Experientia* **17**, 168-169.
- FONG, G.; WALKER, M. A.; GRANETT, J.; 1995: RAPD assessment of California phylloxera diversity. *Mol. Ecol.* **4**, 459-464.
- FORNECK, A.; WALKER, M. A.; BLAICH, R.; 1998: Estimating the genetic diversity among European collections of grape phylloxera (*Daktulosphaira vitifoliae* FITCH) with AFLP-PCR marker. VII<sup>th</sup> International Symposium on Grapevine Genetics and Breeding, 6.-10.6.1998, Montpellier, France.
- GRANETT, J.; GOHEEN, A. C.; LIDER, L. A.; WHITE, J. J.; 1987: Evaluation of grape rootstocks for resistance to type A and type B grape phylloxera. *Amer. J. Enol. Viticult.* **38** (4), 298-230.
- LIN, H.; DOWNIE, D. A.; WALKER, M. A.; GRANETT, J.; ENGLISH-LOEB, G.; 1999: Genetic structure in native populations of grape phylloxera (*Daktulosphaira vitifoliae* FITCH). *Ann. Entomol.* (in press).
- MAILLET, P.; 1957 a: Contribution à l'Étude de la Biologie du Phylloxéra de la Vigne. 1ère éd., Vol. 11. Masson et Coie, Editeurs, Paris.
- -; 1957 b: Sur les chromosomes de quelques phylloxérides de France. *Vitis* **1**, 153-155.
- MANICARDI, G. C.; BIZZARO, D.; AZZONI, P.; BIANCHI, U.; 1997: Cytological and electrophoretic analysis of DNA methylation in the holocentric chromosomes of *Megoura viciae* (Homoptera, Aphididae). *Genome* **37**, 625-630.

Received May 31, 1999