

- MUELLER-DOMBOIS, D.; ELLENBERG, H.; 1974: Aims and Methods of Vegetation Ecology, 84-90. Ed. John Wiley & Sons, N.Y.
- MURISIER, R.; RYSER, J. P.; AERNY, J.; 1982: Fumure potassique et acidité des moûts. Rev. Suisse Viticolt. Arboricolt. Horticult. 14, 33-36.
- OUGH, C. S.; LIDER, L. A.; COOK, J. A.; 1968: Rootstock-scion interactions concerning wine making. I. Juice composition changes and effects on fermentation rate with St. George and 99-R rootstocks at two nitrogen fertilizer levels. Amer. J. Enol. Viticul. 19, 213-227.
- RIBÉREAU-GAYON, J.; PEYNAUD, E.; SUDRAUD, P.; RIBÉREAU-GAYON, P.; 1972: Sciences et Techniques du Vin. Ed. Dunod, Paris.
- SALGUES, M.; 1977: La pratique de la dégustation du vin et son interprétation statistique. Progr. Agric. Vitic. 94, 78-87, 112-123, 137-139.
- SCIENZA, A.; CASASSA, M. T.; VISAI, C.; CONCA, E.; 1984: Il controllo genetico della nutrizione potassica nella vite. Riv. Viticolt. Enol. 37, 682-691.
- SMART, R. E.; 1982: Vine manipulation to improve wine grape quality. Proc. Symp. Grape Wine Cent. Davis, Ca, USA, 362-375.
- ; 1985: Principles of grapevine canopy microclimate manipulation with implications for yield and quality. A review. Amer. J. Enol. Viticul. 36, 230-239.
- ; ROBINSON, J. B.; DUE, G. R.; BRIEN, C. J.; 1985: Canopy microclimate modification for the cultivar Shiraz. I. Definition of canopy microclimate. Vitis 24, 17-31.
- SMITH, R. J.; KLIWER, W. M.; 1984: Estimation of Thompson Seedless grapevine leaf area. Amer. J. Enol. Viticul. 35, 16-22.
- SUDRAUD, P.; 1958: Interprétation des courbes d'absorption des vins rouges. Ann. Techn. Agric. 7, 203-208.
- VIDAL, M.; BLOUIN, J.; 1978: Dosage colorimétrique rapide de l'acide tartrique dans les moûts et les vins. Rev. Fr. Oenol. 16, 39-46.
- WOLPERT, J. A.; HOWELL, G. S.; MANSFIELD, T. K.; 1983: Sampling Vidal Blanc grapes. I. Effect of training systems, pruning severity, shoot exposure, shoot origin, and cluster thinning on cluster weight and fruit quality. Amer. J. Enol. Viticul. 34, 72-76.

Clonal selection of Cabernet

L. BEREZNÁI, Z. BEREZNÁI and L. OLÁH

Egervin Research Centre, P.O.Box 83, H-3301 Eger, Hungary

A b s t r a c t : As a consequence of ecological conditions, in Hungary the grape varieties Cabernet franc and Cabernet Sauvignon became deteriorated and were supplanted by other varieties in production. This was the reason for starting clonal selection of these varieties in Eger in 1962 by the method of four-stage clone selection of M. NEMETH.

As a result of this selection, we produced clones of the two varieties on the basis of 20 years of observation. The clones of Cabernet franc E. 11 and Cabernet Sauvignon E. 153 yield 30-40 % more than the base variety and – depending on the year – they produce wine of excellent or good quality.